

**A
M
I
T**

**ARCHÄOLOGISCHE
MITTEILUNGEN AUS
IRAN UND
TURAN**

**Band 48
2016**

**Exploiting Mangroves - Environmental
changes and human interference along the
northern coast of Arabian Sea during
Holocene**

Exploiting mangroves: Environmental changes and human interference along the northern coast of the Arabian Sea (Pakistan) during the Holocene

By Paolo Biagi, Renato Nisbet and Tiziano Fantuzzi

Schlagwörter: Arabisches Meer, Pakistan, Mangroven, Holozän, Kjökkenmöddinger, ^{14}C -Datierung
Keywords: Arabian Sea, Pakistan, mangroves, Holocene, shell-middens, radiocarbon chronology
Ключевые слова: Аравийское море, Пакистан, мангровые заросли, Голоцен, кьёккенмёддинги, радиоуглерод хронология

Introduction

The scope of this paper is to overview our knowledge of the prehistory of the northern coast of the Arabian Sea in Lower Sindh and Las Bela in Balochistan, define the chronology of the earliest coastal settlements, and discuss their location, characteristics, and disappearance. Coastal archaeology has greatly improved during the last 30 years¹ thanks to the interest that many archaeologists, palaeoclimatologists and geomorphologists, have paid to this unique field of research. This fact led to the discovery of the earliest traces of coastal navigation,² whose archaeological evidence, in the study region, is marked by the impressive finds excavated at as-Sabiyah in Kuwait, at the westernmost edge of the Persian/Arabian Gulf.³

Almost nothing was known of the presence of archaeological sites along the northern coast of the Arabian Sea until the second half of the 1970s. In those years Professor A. R. Khan of the Department of Geography, Karachi University, carried out a systematic geoarchaeological survey of the entire territory around Karachi. Thanks to his discoveries at present we know that the whole area is very rich in sites of different ages, spanning a long period, from the Middle Palaeolithic to the Buddhist period.⁴

Unfortunately most of the sites discovered by Professor A. R. Khan are still unpublished, and too many of his important discoveries are unknown or unreported by most archaeologists.⁵ However, our knowledge of the archaeology of the northern part of the Arabian Sea has dramatically improved following the results achieved by his surveys.

It is again thanks to the work of the aforementioned author that the Neolithic and Bronze Age shell middens of the Bay of Daun were discovered

in Las Bela in 2000. The sites were systematically surveyed in 2004 and 2008.⁶ Between 2000 and 2014 a few visits were paid also to Capes Gadani and Phuari,⁷ Rehri, along the coastal terraces that extend east of Karachi, the Tharro Hills,⁸ Balakot,⁹ Pir Shah Jurio,¹⁰ and Sonari near Cape Monze (Ras Mauri).¹¹

Even more surprising is that the surveys carried out by other authors before the 1970s along the same coastline as far as Makran, did not yield any trace of the most characteristic sites of the Arabian Sea coast, namely shell middens.¹² This evidence strangely contrasts with the data available from the Sultanate of Oman since the 1960s, and more generally along the western coast of the Arabian Sea.¹³

The shell middens of the Bay of Daun were discovered in January 2000 during a short visit paid to the coast of Las Bela together with Professor A. R. Khan. The main scope of the visit was to collect complete specimens of *Terebralia palustris* mangrove shells, a few samples of which Professor Khan kept in the stores of the Museum of Prehistory and Palaeogeography, Institute of Geography, Karachi University.

Following the discovery of the Daun shell middens, the surveys were extended first to the Indus Delta, in Lower Sindh,¹⁴ and later to the coast of Las Bela, in Balochistan.¹⁵ Both areas yielded an impressive number of archaeological sites, mainly *Terebralia palustris* and *Telescopium telescopium* shell middens, and shell scatters both close to the seaside and well inland. A good series of AMS dates was obtained from these sites, whose scope was

¹ Bailey/Parkington 1988; Erlandson/Fitzpatrick 2006; Rainbird 2007; Erlandson/Rick 2008.

² Bjerck 2017.

³ Carter/Crawford 2010.

⁴ Khan 1979a.

⁵ Wright 2010; Coningham/Young 2015; Kenoyer 2015.

⁶ Biagi et al. 2012.

⁷ Biagi et al. 2013a.

⁸ Majumdar 1934; Piggott 1950; Fairservis 1975.

⁹ Khan 1979c, 75.

¹⁰ Khan 1979a, 3.

¹¹ Biagi/Nisbet 2014.

¹² Stein 1943; Snead 1966; 1969; Dales 1982; Besenval 1992;

Dales/Lipo 1992; Hasan 2002.

¹³ Biagi 1988; Cleuziou 2004; Berger et al. 2005; Амирханов

2006.

¹⁴ Biagi 2010.

¹⁵ Biagi 2011; Biagi et al. 2013b.

to achieve a first reliable sequence of the prehistoric settlement and environmental changes that took place in this almost forgotten region of the north Arabian Sea coast since the beginning of the Holocene.

Current and historical mangroves in Pakistan

It is well known that mangroves represent an exceptional environment, perfectly adapted to inter-tidal habitat such as estuaries and deltas.¹⁶ They are so strictly connected to the coast that some authors define mangroves to include the formation below the high tide mark (*tidal forests*), though other mangroves “may occur from far below the level of the lowest to above the level of the highest tides, or on coasts where there are no tides at all”.¹⁷ The fact that they live in areas where both fresh and saline water occur, favour the reconstruction of the coastal variations in those regions, the Indus Delta for instance, where dramatic changes in the landscape, not yet fully understood, took place during the Holocene.¹⁸

Many authors have pointed out the economic importance of mangroves, providing fuelwood,¹⁹ timber, tannin and medicinal products as well as crustaceans and fish for food,²⁰ though no recent gatherings of the most frequent, typical molluscs have ever been mentioned,²¹ as happened in prehistoric times in the Gulf and along the coasts of the Arabian Sea.²²

Fig. 1 Distribution map of the present mangroves along the coasts of Sindh and Balochistan (Pakistan) (topographic map OpenStreet Map; drawing by R. Nisbet)



- ¹⁶ Kathiresan/Rajendran 2005; Schwadron 2013.
¹⁷ Lugo/Snedaker 1974, 43.
¹⁸ Pithawalla 1939; 1976; Wilhelmy 1968, 1986; Flam 1987, 1999; Kevran 1999.
¹⁹ Aitken 1907, 32; Hasan 2002, 10.
²⁰ Tomlison 1986; Hogarth 1999; Ellison 2014.
²¹ Siddiqi 1956; Haas 1959.
²² Biagi 2008; Boivin/Fuller 2009.

Out of Pakistan’s 1000 km coastline, roughly 300 are covered by mangroves. In the case of the Indus Delta, these tidal forests border the numerous creeks extending well inland, at a distance of 30 km or more from the coast, covering a huge territory of ca. 600,000 ha.²³ A much lesser extent have the other four mangrove areas, found at Manora (Karachi),²⁴ Miāni Hor,²⁵ Kalamat Khor and Jiwani (the last three in Balochistan), together forming only 5% of the total Pakistani mangroves endangered by increasing human pressure²⁶ (Fig. 1).

Ancient historians have left some descriptions of mangroves at the borders of the Indian Ocean and the Gulf. Probably the earliest mention is found in Theophrastus who, in his *Historia Plantarum* (305 BC), reports from Aristobulos on the voyage by Nearchos in 325 BC from the Indus Delta to the Persian Gulf. The trees described by Alexander’s admiral would apply to *Rhizophora* sp. and *Avicennia* sp.²⁷ Arrian mentions of mangroves (“These trees were on ground which was left dry by the ebb-tide; but when the water advanced they looked as if they had grown in the sea”: Arrian, *Anabasis*, VI.22.6)²⁸ in his description of Alexander’s voyage are the last one together with that of Pliny’s *Naturalis Historia*, XIII.25.51.²⁹ Apart from early short points in Theophrastus and Strabo, before the 13th century AD observations are reported by the botanist and physician Ibn al-Bitār along the Arabian coasts.³⁰

According to B. Rollet³¹ in the following three centuries no more than five or six descriptions of mangroves in the world botanical literature are found before van Rheedee’s *Hortus malabaricus*,³² broadly referring to Indian Ocean mangroves. Paradoxically, Aitchison’s list of Punjab and Sindh plants reports the presence of *Rhizophora* sp. at the Indus Delta, but not of *Avicennia marina*, by far more common.³³

Even the otherwise always very well informed *Gazetteer of Sind*³⁴ pays no attention to the coastal vegetation between Karachi and the Indus Delta. More details were provided by A. J. Murray (1881), who described some mangrove trees (*Ceriops candoleana*) as common “at the mouth of Indus, and in the salt-water creeks, Kurrachee, and on the coast”,³⁵ which is certainly an overestimation of

- ²³ Snedaker 1984, 256; Qureshi 1990.
²⁴ Baillie 1890, 52-53; Campbell 1999.
²⁵ Saifullah et al. 2002.
²⁶ Hameed-Baloch et al. 2014.
²⁷ Eggermont 1975.
²⁸ Ambaglio 1994, 561.
²⁹ McCrindle 1901; 1972.
³⁰ Ibn al-Bitār 1987.
³¹ Rollet 1981.
³² van Rheedee 1678-1703.
³³ Aitchison 1869.
³⁴ Hughes 1876.
³⁵ Murray 1881, 190.

the occurrence of this species at present known as *Ceriops tagal* (Perr.) C. B. Rob., and also *Rhizophora*, *Bruguiera* and *Avicennia*. More precise indications on the location of mangroves in the Indus Delta are found in the Gazetteer of the Province of Sind, which mentions three species of mangrove trees,³⁶ though the first large account on Indus mangroves will appear only several years later.³⁷

Balochistan mangroves have attracted attention much later, since 1980, because of their potential economic interest, and important protective programmes, while repopulation projects have been since carried out.³⁸ Moreover, Miāni Hor is the only area in Pakistan where the three mangrove species *Avicennia marina*, *Rhizophora mucronata* and *Ceriops tagal* occur naturally.³⁹

Both Sindh and Balochistan mangroves urgently need long-term rehabilitation projects, as widely recognised by local and international authorities, in order to stop the erosion of their fragile ecosystems, whose overall extension changed from over 600,000 ha in 1932 to ca. 85,000 ha in 2005⁴⁰ mostly due to progressively increasing human pressure and pollution.⁴¹

Methodology

The use of Gastropods as palaeoenvironmental indicators of mangroves is well known.⁴² Their exploitation is well recorded from many archaeological sites of the Gulf of Oman and Yemen along the western coast of the Arabian Sea,⁴³ the Persian/Arabian Gulf,⁴⁴ and the Red Sea.⁴⁵ The technique is based on the close relationship between the mangrove unique environment and a few species of molluscs (in particular *Terebralia palustris* L. and *Telescopium telescopium* L.), which feed mostly on fallen leaves of *Rhizophora* and *Avicennia*, living under the thick mangrove canopy as well as in the surrounding more open, muddy areas⁴⁶ (Fig. 2).

The frequent exploitation of these molluscs as food by ancient groups led to the formation of characteristic low mounds, or sometime simply



Fig. 2 Miāni Hor, south-easternmost creek: *Rhizophora* sp. and *Avicennia* sp. mangrove forest (top), and *Telescopium telescopium* gastropods (bottom) (Photographs by P. Biagi and R. Nisbet 2012)

scatters of debris. The shells, which were in a few cases minutely fragmented in order to extract the edible soft tissues, are usually found concentrated in areas ranging from small spots 2-3 m diameter to very large areas, up to more than 40 m diameter (shell middens) (Fig. 3). The identification of shell concentrations was generally made easier due to the barren nature of the surface. Chipped stone artefacts (mostly flint/chert) in close relationship with the shells were made using, at least in some cases, local outcrops (as, for example, at Siranda and other coastal sites in Las Bela, reddish-brown flint from Cape Gadani).⁴⁷

The shells primarily consist of typical mangrove gastropods (*T. palustris* and *T. telescopium*) and bivalves (mostly *Anadara rhombea* Born, *Anadara cf. uropygmelana*, *Circenita callipyga* Born, *Thais lacera* Born etc.) distinctive of both mangroves and shallow waters of the upper intertidal zone, on sandy and muddy bottoms.⁴⁸ Lists of mol-

³⁶ Aitken 1907, 31.

³⁷ Blatter et al. 1929.

³⁸ Rasool/Saifullah 2005; Amjad/Kamaruzaman 2007.

³⁹ Baig/Iftikhar 2006, 6.

⁴⁰ Damhoureyeh/Ghalib 2014.

⁴¹ Rizwi et al. 1999; Amjad et al. 2007; Hameed-Baloch et al. 2014.

⁴² Thomas 2015a; 2015b.

⁴³ Biagi/Nisbet 1992; 2006; Plaziat 1995; Амирханов 1997; Amirkhanov et al. 2001; Uerpman/Uerpman 2003; Berger et al. 2013.

⁴⁴ Choquert 1980; Feulner 2006; Hellyer/Aspinal 2006.

⁴⁵ Cattani/Bökönyi 2002; Bar-Yosef Mayer/Beyn 2009, 2013; Bailey et al. 2013.

⁴⁶ Vannini et al. 2008.

⁴⁷ Biagi et al. 2013a Plate 1.

⁴⁸ Biagi et al. 2013c.



Fig. 3 Lake Siranda: Shell midden 29 (SRN-29), main site (top), and small heap of fragmented mangrove and marine shells east of the same site (bottom) (photographs by P. Biagi 2012)



Fig. 4 Balakot: The archaeological site (BLK-1) from the south (photograph by P. Biagi 2012)

luscs found in our surveys have been provided in previous papers.⁴⁹

Palynological evidence of palaeo-mangroves in Pakistan is very limited. Margaret McKean,⁵⁰ in her study on the ancient vegetation of Balakot⁵¹ (Fig. 4) did not identify any pollen of *Avicennia* or *Rhizophora*, in spite of the proximity of the archaeological site to the well-documented Siranda mangroves during the same Bronze Age centuries. In a more recent work on a marine core off the Makran coast, Ivory and Lézine⁵² described, amongst other continental vegetation associations, a well-defined curve of mangrove pollen types, with its maximum value starting from ca. the mid 6th millennium BP. The palaeoclimatic interpretation of the diagram points to humid conditions that favoured the diffusion of *Rhizophora* and freshwater algae until the following millennium, when the humid Holocene Atlantic period came to an end, and gradually more xeric vegetation indicators prevailed.

A very different picture can be drawn from the numerous, well-documented palynological sites along the western coast of the Indian Subcontinent.⁵³ In this latter region, under quite a different climatic regime,⁵⁴ mangroves were diffused before 40 ka cal BP and lasted until the mid-Holocene, when the

⁴⁹ Biagi et al. 2013a, 2013c.

⁵⁰ McKean 1983.

⁵¹ Dales 1979; Shaffer 1986.

⁵² Ivory/Lézine 2009.

⁵³ Kumaran et al. 2005.

⁵⁴ Gupta 2004.

weakening of the monsoon led gradually to drying up most of the coastal vegetation.⁵⁵

Until now no macro-remains (fruits, charred wood etc) from palaeo-mangroves have been recognised in Pakistan, in spite of R. E. Snead's claim⁵⁶ of the presence of mangrove along the western border of Lake Siranda as late as 1890. Careful inspections carried out in the area in 2012 and 2013 did not yield any evidence of mangrove trees or faunal remains, which have totally disappeared from Siranda ca. 4500 years, following the progressive drying up of the basin during the Bronze Age.

The research area

The archaeological surveys carried out in Lower Sindh and Las Bela (Balochistan) between 2000 and 2014 led to the discovery of 17 locations with molluscs evidence of palaeo-mangroves, close to the present coastline and also far inland, from Miāni Hor (Las Bela), to the Makli Hills (Thatta, Sindh). With the present 84 radiocarbon dates from *Terebralia palustris* and *Telescopium telescopium*⁵⁷ (Table 1), on which the present paper is mostly based, and 22 more from marine shells (*Purpura panama*, *Meretrix* sp., *Ostreidae*, *Lunella coronata*, *Turbo bruneus*, *Mactridae*), the project still underway furnishes the first detailed data-set for the reconstruction of the early exploitation (second half of the 9th millennium BP) of mangal resources by semi-nomadic populations. Furthermore it adds new arguments to the interpretation of the evolution of the prograding Indus Delta during the last eight millennia.

⁵⁵ Gupta et al. 2003; Yoganandan et al. 2013; Zorzi et al. 2015.

⁵⁶ Snead 1966 Fig. 21, quoting the Service of India Edition 1915.

⁵⁷ Reid et al. 2008.

Table 1 List of the radiocarbon dates from mangrove gastropods from the sites mentioned in the text. Calibrations according to the marine curve by Reimer et al. 2013

Site name	Coordinates	Altitude (m)	Material	Lab. n°	δ ¹³ C	Uncal BP	Cal BC 2σ	Reference
SRN-43 (Lake Siranda - Las Bela, Balochistan)	25°30'25.3"N-66°38'31.7"E	8	T. palustris	GrA-54290	-3,55	7200±35	5610-5453	Biagi 2013
SRN-38 (Lake Siranda - Las Bela, Balochistan)	25°30'07.0"N-66°38'44.7"E	9	T. palustris	GrA-54303	-6,58	7095±35	5517-5332	Biagi 2013
SRN-56 (Lake Siranda - Las Bela, Balochistan)	25°29'56.3"N-66°38'56.9"E	10	T. palustris	GrA-57702	-6,17	6980±35	5436-5229	Unpublished
SRN-33 (Lake Siranda - Las Bela, Balochistan)	25°29'58.4"N-66°39'16.0"E	12	T. palustris	GrA-54291	-6,16	6770±35	5222-4978	Biagi 2013
SRN-32 (Lake Siranda - Las Bela, Balochistan)	25°29'59.5"N-66°39'17.1"E	12	T. palustris	GrA-57528	-6,66	6630±35	5049-4796	Unpublished
SRN-37 (Lake Siranda - Las Bela, Balochistan)	25°29'59.3"N-66°38'57.3"E	7	T. palustris	GrA-55821	-5,87	6595±45	5026-4746	Biagi et al. 2013a
SRN-29 (Lake Siranda - Las Bela, Balochistan)	25°30'26.8"N-66°37'35.1"E	10	T. palustris	GrA-54299	-5,57	6595±35	5008-4764	Biagi 2013
SRN-66 (Lake Siranda - Las Bela, Balochistan)	25°30'51.8"N-66°36'52.9"E	8	T. palustris	GrA-57703	-5,27	6575±35	4984-4740	Unpublished
SRN-64 (Lake Siranda - Las Bela, Balochistan)	25°31'18.0"N-66°36'43.2"E	13	T. palustris	GrA-57535	-5,19	6515±35	4917-4683	Unpublished
SRN-28.10 (Lake Siranda - Las Bela, Balochistan)	25°30'30.6"N-66°37'35.4"E	16	T. palustris	GrA-62260	-4,78	6500±40	4914-4659	Unpublished
SRN-67 (Lake Siranda - Las Bela, Balochistan)	25°30'43.8"N-66°36'52.8"E	11	T. palustris	GrA-59841	-4,75	6370±60	4780-4461	Unpublished
SRN-39bis (Lake Siranda - Las Bela, Balochistan)	25°30'08.5"N-66°38'41.2"E	9	T. telescopium	GrA-54298	-4,53	6335±35	4696-4467	Biagi 2013
SRN-63.2 (Lake Siranda - Las Bela, Balochistan)	25°32'31.1"N-66°37'09.5"E	7	T. palustris	GrA-57534	-4,1	6325±35	4686-4456	Unpublished
SRN-1 (Lake Siranda - Las Bela, Balochistan)	25°31'19.3"N-66°36'39.6"E	5	T. palustris	GrA-50325	-6,213	6305±40	4682-4436	Biagi 2013
SRN-62 (Lake Siranda - Las Bela, Balochistan)	25°31'28.8"N-66°36'44.4"E	5	T. palustris	GrA-59842	-4,73	6230±60	4630-4328	Unpublished
SRN-75 (Lake Siranda - Las Bela, Balochistan)	25°32'29"N-66°37'15"E	5	T. palustris	GrA-63864	-6,8	6220±40	4572-4338	Unpublished

Site name	Coordinates	Altitude (m)	Material	Lab. n°	$\delta^{13}C$	Uncal BP	Cal BC 2 σ	Reference
SRN-40 (Lake Siranda - Las Bela, Balochistan)	25°30'09.9"N-66°38'40.4"E	4	T. palustris	GrA-55823	-3,86	6145±45	4496-4270	Biagi et al. 2013a
SRN-39 (Lake Siranda - Las Bela, Balochistan)	25°30'08.2"N-66°38'41.4"E	9	T. telescopium	GrA-55822	-4,33	6145±45	4496-4270	Biagi et al. 2013a
SRN-76 (Lake Siranda - Las Bela, Balochistan)	25°32'20"N-66°37'07"E	5	T. palustris	GrA-59840	-3,64	6100±60	4488-4212	Unpublished
SRN-63 (Lake Siranda - Las Bela, Balochistan)	25°31'19.3"N-66°36'39.4"E	6	T. palustris	GrA-63868	-4,01	6055±40	4419-4195	Unpublished
SRN-2 (Lake Siranda - Las Bela, Balochistan)	25°31'31.0"N-66°36'48.9"E	0	T. palustris	GrA-50323	-4,638	5950±40	4306-4045	Biagi et al. 2013a
SRN-31 (Lake Siranda - Las Bela, Balochistan)	25°30'01.1"N-66°39'19.0"E	4	T. palustris	GrA-55820	-5,03	5875±45	4230-3967	Biagi et al. 2013a
SRN-47 (Lake Siranda - Las Bela, Balochistan)	25°30'39.9"N-66°38'06.3"E	10	T. palustris	GrA-54296	-3,46	5800±35	4154-3920	Biagi et al. 2013a
SRN-23 (Lake Siranda - Las Bela, Balochistan)	25°30'47.7"N-66°37'39.2"E	7	T. palustris	GrA-54294	-4,67	5780±30	4118-3900	Biagi et al. 2013a
SRN-42 (Lake Siranda - Las Bela, Balochistan)	25°30'25.1"N-66°38'32.2"E	11	T. palustris	GrA-54292	-5,79	5755±35	4070-3810	Biagi et al. 2013a
SRN-73 (Lake Siranda - Las Bela, Balochistan)	25°30'26.8"N-66°37'31.7"E	9	T. palustris	GrA-57707	-3,9	5695±35	3988-3770	Unpublished
SRN-44 (Lake Siranda - Las Bela, Balochistan)	25°30'22"N-66°38'38"E	2	T. palustris	GrA-54301	-7,2	5690±35	3982-3766	Biagi et al. 2013a
SRN-24 (Lake Siranda - Las Bela, Balochistan)	25°30'48.0"N-66°37'37.4"E	5	T. telescopium	GrA-55818	-6,12	5665±45	3970-3718	Biagi et al. 2013a
SRN-72 (Lake Siranda - Las Bela, Balochistan)	25°29'31.9"N-66°36'54.7"E	7	T. palustris	GrA-57704	-4,67	5665±35	3961-3745	Unpublished
SRN-52 (Lake Siranda - Las Bela, Balochistan)	25°30'39.9"N-66°38'13.0"E	16	T. palustris	GrA-57701	-5,61	5575±35	3894-3650	Unpublished
SRN-28 (Lake Siranda - Las Bela, Balochistan)	25°30'32.4"N-66°37'35.9"E	9	T. palustris.	GrA-55819	-2,55	5440±40	3726-3510	Biagi et al. 2013a
SRN-16 (Lake Siranda - Las Bela, Balochistan)	25°31'39.1"N-66°35'53.9"E	7	T. palustris	GrA-55817	-3,86	5065±40	3329-3023	Biagi et al. 2013a
SRN-57 (Lake Siranda - Las Bela, Balochistan)	25°31'32.0"N-66°36'48.7"E	11	T. palustris	GrA-57533	-0,55	4315±35	2326-2026	Unpublished
BLK-1 (Balakot - Las Bela, Balochistan)	25°27'22.6"N-66°42'39.7"E	27	T. palustris	GrA-55828	-3,77	4660±40	2831-2498	Biagi et al. 2013a
Daun-110 (Daun Bay - Las Bela, Balochistan)	25°00'00"N-66°42'21"E	7	T. palustris	GrN-31492	-3,44	6690±40	5176-4875	Biagi 2011
Daun-111 (Daun Bay - Las Bela, Balochistan)	24°59'59"N-66°42'25"E	9	T. palustris	GrN-31493	-3,57	6590±45	5019-4739	Biagi 2011
Daun-1 (Daun Bay - Las Bela, Balochistan)	25°00'15"N-66°42'39"E	9	T. palustris	GrN-26368	-3,08	6380±40	4762-4506	Biagi 2004
Daun-10 (Daun Bay - Las Bela, Balochistan)	25°00'13"N-66°42'45"E	8	T. palustris	GrN-31489	-3,97	6305±45	4690-4427	Biagi 2011
Daun-6 (Daun Bay - Las Bela, Balochistan)	24°59'20"N-66°42'31"E	19	T. palustris	GrN-28802	1,27	5370±35	3650-3439	Biagi 2011
Daun-116 (Daun Bay - Las Bela, Balochistan)	25°00'07.9"N-66°42'23.7"E	7	T. palustris	GrA-66637	-3,52	5360±40	3639-3384	Unpublished
Daun-5 (Daun Bay - Las Bela, Balochistan)	24°59'19"N-66°42'29"E	19	T. palustris	GrN-28801	-5,44	4900±35	3072-2860	Biagi 2004
Daun-112 (Daun Bay - Las Bela, Balochistan)	25°00'00"N-66°42'28"E	16	T. palustris	GrN-32462	-4,95	4625±30	2748-2464	Biagi 2011
Daun-102 (Daun Bay - Las Bela, Balochistan)	24°59'37"N-66°42'19"E	10	T. palustris	GrN-32117	-5,96	4590±35	2702-2431	Biagi 2011
Daun-105 (Daun Bay - Las Bela, Balochistan)	24°59'34"N-66°42'21"E	9	T. telescopium	GrN-31643	-5,09	4470±40	2546-2247	Biagi 2011
Daun-104 (Daun Bay - Las Bela, Balochistan)	24°59'37"N-66°42'19"E	10	T. palustris	GrN-32118	-6,1	4470±35	2540-2261	Biagi 2011
Daun-101 (Daun Bay - Las Bela, Balochistan)	24°59'37"N-66°42'19"E	10	T. palustris	GrN-31490	-5,49	4470±30	2528-2266	Biagi 2011

Site name	Coordinates	Altitude (m)	Material	Lab. n°	$\delta^{13}\text{C}$	Uncal BP	Cal BC 2 σ	Reference
Daun-113 (Daun Bay - Las Bela, Balochistan)	25°00'03"N-66°42'22"E	7	T. palustris	GrN-32463	-5,44	4455±30	2486-2230	Biagi 2011
Daun-103 (Daun Bay - Las Bela, Balochistan)	24°59'35"N-60°42'22"E	9	T. palustris	GrN-31491	-5,37	4435±40	2475-2192	Biagi 2011
Daun-119 (Daun Bay - Las Bela, Balochistan)	25°00'25"N-66°43'06"E	6	T. palustris	GrN-31644	-4,05	4165±25	2106-1870	Biagi 2011
Daun-3 (Daun Bay - Las Bela, Balochistan)	25°00'26"N-66°43'04"E	4	T. palustris	GrN-27945	-4,49	4100±30	2010-1752	Biagi 2004
Daun-117 (Daun Bay - Las Bela, Balochistan)	25°00'07"N-66°42'22"E	7	T. palustris	GrN-31494	-3,95	1440±30	1086-1282 AD	Unpublished
GDN-0 (Cape Gadani - Las Bela, Balochistan)	25°06'42.4"N-66°43'13.2"E	24	T. palustris	GrN-26369	-4,99	4460±30	2494-2234	Biagi 2004
PHR-11 (Ras Phuari - Las Bela, Balochistan)	25°05'19.0"N-66°42'26.93"E	19	T. palustris	GrA-55826	-5,09	4415±40	2467-2186	Biagi et al. 2013b
SNR-101 (Sonari, Hab River Mouth - Sindh)	24°52'38.7"N-66°41'46.78"E	12	T. palustris	GrA-62252	-4,2	4690±35	2844-2564	Unpublished
SNR-7 (Sonari, Hab River Mouth - Sindh)	24°52'27.7"N-66°41'37.8"E	14	T. palustris	GrA-59832	-2,36	4560±60	2734-2316	Unpublished
SNR-4bis/1 (Sonari, Hab River Mouth - Sindh)	24°52'39.4"N-66°41'35.2"E	27	T. palustris	GrA-62250	-3,79	4520±35	2586-2313	Unpublished
SNR-5 (Sonari, Hab River Mouth - Sindh)	24°52'38.3"N-66°41'34.9"E	27	T. telescopium	GrA-59833	-5,14	4470±60	2566-2196	Biagi and Nisbet 2014
SNR-8 (Sonari, Hab River Mouth - Sindh)	24°52'13.5"N-66°41'18.4"E	23	T. palustris	GrA-62251	-4,38	4405±35	2452-2176	Unpublished
Sonari (Sonari, Hab River Mouth - Sindh)	24°52'28"N-66°41'54"E	27	T. palustris	GrN-27054	-4,43	4080±30	1986-1731	Biagi 2004
SNR-4bis/2 (Sonari, Hab River Mouth - Sindh)	24°52'38.8"N-66°41'34.6"E	24	T. palustris	GrA-66633	-7,47	3995±35	1879-1627	Unpublished
SNR-1D (Sonari, Hab River Mouth - Sindh)	24°52'37.5"N-66°41'31.7"E	27	T. telescopium	GrA-59835	-4,42	3660±50	1491-1204	Biagi and Nisbet 2014
SNR-3 (Sonari, Hab River Mouth - Sindh)	24°52'38.2"N-66°41'41.4"E	9	T. palustris	GrA-62249	0,94	2190±30	329-560 AD	Unpublished
SNR-2 (Sonari, Hab River Mouth - Sindh)	24°52'38.9"N-66°42'02.6"E	3	T. telescopium	GrA-59834	-5,1	670±50	1800- AD	Unpublished
PSH-1bis (Pir Shah Jurio, Hab River Mouth - Sindh)	24°55'39.1N-66°44'28.2"E	35	T. palustris	GrA-66638	-4,15	4270±35	2265-1974	Unpublished
PSH-1 (Pir Shah Jurio, Hab River Mouth - Sindh)	24°55'39.1N-66°44'28.2"E	35	T. palustris	GrN-26370	-4,38	4130±20	2035-1806	Biagi 2004
RHR-3bis (Rehri, Karachi - Sindh)	24°49'12"N-67°13'42"E	10	T. palustris	GrA-66631	-4,13	7045±45	5483-5287	Unpublished
MH-15 (Mulri Hills, Karachi - Sindh)	24°55'41"N-67°07'14"E	67	T. palustris	GrA-63863	-4,01	7320±40	5711-5524	Unpublished
MH-14 (Mulri Hills, Karachi - Sindh)	24°55'42"N-67°07'25"E	65	T. telescopium	GrA-63869	-4,57	6155±40	4504-4300	Unpublished
MH-4B (Mulri Hills, Karachi - Sindh)	24°55'47"N-67°07'57"E	65	T. palustris	GrA-66630	-5,24	6035±40	4379-4156	Unpublished
MH-18 (Mulri Hills, Karachi - Sindh)	24°54'45"N-67°06'30"E	65	T. palustris	GrA-23639	-6,6	5790±70	4211-3816	Biagi 2004
MH-17 (Mulri Hills, Karachi - Sindh)	24°54'43"N-67°07'55"E	65	T. palustris	GrA-66634	-3,98	5530±40	3850-3617	Unpublished
Garo-1 (Bhambor - Sindh)	24°45'36.3"N-67°33'17.4"E	31	T. telescopium	GrA-59844	-3,64	6320±60	4726-4408	Unpublished
THR-3 (Tharro Hills, Gujo - Sindh)	24°43'46"N-67°45'07"E	13	T. palustris	GrA-47084	-5,15	5555±35	3876-3635	Biagi 2011
Beri (Gujo - Sindh)	24°43'00"N-67°45'09"E	7	T. palustris	GrN-32166	-6,9	5960±50	4320-4041	Biagi 2010
JSH-1bis (Shah Husein, Gujo - Sindh)	24°42'26.0"N-67°48'38.3"E	12	T. telescopium	GrA-66636	-4,79	5800±40	4165-3910	Unpublished
JSH-2 (Shah Husein, Gujo - Sindh)	24°42'26"N-67°48'39"E	19	T. telescopium	GrA-45181	-3,21	4245±40	2230-1926	Biagi 2010

Site name	Coordinates	Altitude (m)	Material	Lab. n°	$\delta^{13}C$	Uncal BP	Cal BC 2 σ	Reference
JSH-10 (Shah Husein, Gujo - Sindh)	24°42'09.8"N-67°48'28.1"E	14	T. telescopium	GrA-62255	-5,18	2715±30	339-78	Unpublished
KKT-2 (Kalan Kot, Thatta - Sindh)	24°42'17.3"N-67°52'23.5"E	22	T. palustris	GrN-32464	-5,5	6320±45	4700-4442	Biagi 2011
MKL-10 (Makli Hills, Thatta - Sindh)	24°37'40.6"N-67°51'41.2"E	25	T. telescopium	GrA-62256	-7,02	6140±40	4486-4274	Unpublished
MKL-1 (Makli Hills Thatta - Sindh)	24°36'52.5"N-67°51'36.5"E	24	T. palustris	GrA-50330	-3,929	5750±40	4074-3796	Unpublished
KKT-4 (Kalan Kot, Thatta - Sindh)	24°42'15.3"N-67°52'15.7"E	27	T. telescopium	GrA-59843	-7,03	5460±60	3788-3498	Unpublished
KKT-3 (Kalan Kot, Thatta - Sindh)	24°41'54.8"N-67°52'40.4"E	32	T. telescopium	GrA-50324	-5,01	5270±40	3579-3336	Unpublished
OBS-1 (Aban Shah, Thatta - Sindh)	24°22'17.8"N-67°58'20.6"E	8	T. palustris	GrA-47082	-9,17	3790±35	1616-1398	Biagi 2011
KRM-13 (Kot Raja Manjera, Jerrack - Sindh)	25°01'21"N-68°12'37"E	45	T. palustris	GrA-47083	-6,17	4635±35	2771-2469	Biagi 2011




Fig. 5 Distribution map of the radiocarbon-dated shell middens and shell scatters in the study area: A) 8th millennium BP (light blue): SRN-43 (1), SRN-38 (2), MH-15 (3), RHR-3bis (4); 9th millennium BP: KDJ-1 (5) (black); B) 7th millennium BP (blue): SRN-56, SRN-33, SRN-32, SRN-37, SRN-29, SRN-66, SRN-64, SRN-28.10, SRN-67, SRN-39bis, SRN-63.2, SRN-1, SRN-62, SRN-75, SRN-40, SRN-39, SRN-76, SRN-60 (1-18), Daun-110, Daun-111, Daun-1, Daun-10 (19-22), MH-14 (23), MH-4B (24), GARO-1 (25), KKT-2 (26), MKL-10 (27); C) 6th millennium BP (green): SRN-2, SRN-31, SRN-47, SRN-23, SRN-42, SRN-73, SRN-44, SRN-24, SRN-72, SRN-52, SRN-28, SRN-16 (1-12), Daun-6 (13), Daun-116 (14), MH-18 (15), MH-17 (16), THR-3 (17), Beri (18), JSH-1bis (19), MKL-1 (20), KKT-4 (21), and KKT-3 (22) (drawings by R. Nisbet and P. Biagi)

The surveyed region spans from the shallow depression of Siranda (Las Bela district, Balochistan), the high coastline between Cape Gadani and the mouth of the Hab River, to the flats of the Indus Delta with some remnants of the pre-deltaic Eocene limestone reliefs, the Makli Hills, south of Thatta, and Aban Shah in the lower Delta plain.⁵⁸ Moreover, the research carried out along the limestone mesas of the western side of the Indus, between Ongar, Meting and Jhimpir, while resulting in the discovery of many archaeological sites from the Lower Palaeolithic to Historical times,⁵⁹ yielded evidence of Bronze Age *T. palustris* shells at Kot Raja Manjera, near Jerrack (Jhirak).⁶⁰ This is, at the moment, the northernmost point from which mangrove shells have been found, some 150 km as the crow flies from the present Indus mouth (Fig. 5 and 6).

Before the construction of dams and barrages along its course,⁶¹ the Delta shoreline advanced at an average rate of ca. 45 m/year, with maxima of 150 m/year at the mouth of active channels.⁶² The importance of recognising the complex history of

⁵⁸ Blanford 1880.

⁵⁹ Starnini/Biagi 2011.

⁶⁰ Khan 1979a, 6, 1979c, 71-72; Cousens 1998, 87; Biagi 2010.

⁶¹ Pithawalla 1939; Panhwar 1964; Rahman 1988.

⁶² Hayter 1960; Giosan et al. 2006.

the coastline changes in the millennia,⁶³ in relationship with the prehistoric human presence, is easily understood. Several calculations have been made also regarding the rate of rising of the alluvial plain in the Delta area. According to H. T. Lambrick⁶⁴ a rise of 20 to 30 cm a century is quite a realistic figure, and the central part of the plain would have increased for ca. 9 m in the last 5,000 years. More recent estimates⁶⁵ display offshore sedimentation rates of 50 cm/year at the mouth of the active channel, in the Indus canyon, and huge quantities of sediments, calculated in terms of 250 megatonne per year, were supplied by the river prior to modern damming.⁶⁶

Mangroves and radiocarbon dates

As reported above, the 2000-2014 surveys carried out along the Las Bela coast and the Indus Delta led to the discovery of dozens of prehistoric sites. The research was coupled with a wide programme of radiocarbon dating made mainly on mangrove, but also marine shells, whose scope was to define the chronology of the prehistoric exploitation of the coastal forests. Furthermore carbon isotopes are an excellent way to understand the changing landscape of the Indus Delta, one of the most active coastlines of the world.⁶⁷

The description of the aforementioned changes are based on radiocarbon dates mainly obtained from the apex of one single specimen of adult, decoloured, mangrove Gastropod weighing 5 to 10 grams, collected from the centre of each site. *T. palustris* was preferably selected or, in its absence, *T. telescopium* (see **Table 1**). They all were dated at Groningen University Isotope Laboratory: GrA- in the case of AMS dates from one individual, and more specimens of the same species in the case of ordinary dates (GrN-) processed before 2004. The dates throughout the text are given as uncal BP unless otherwise stated.

Three main large areas have been considered: 1) the Siranda basin, at present a sabkha-like saline depression, some 15 km long and 4 km wide, whose south-western side lies 5 km from the present Sonmiani Lagoon (Miāni Hor) and ca. 15 km from the present sea-shore; 2) the coast between Cape Gadani and the mouth of the Hab River that marks the boundary between Balochistan and Sindh, with its long low-lying shores (like partially at Daun) and some




Fig. 6 Distribution map of the radiocarbon-dated shell middens and shell scatters in the study area: A) 5th millennium BP (yellow): SRN-57 (1), BLK-1 (2), Daun-5, Daun-112, Daun-102, Daun-105, Daun-104, Daun-101, Daun-113, Daun-103, Daun-119, Daun-3 (3-12), GDO-0 (13), PHR-11 (14), SNR-101, SNR-7, SNR-4bis/1, SNR-5, SNR-8, Sonari (15-20), PSH-1bis (21), PSH-1 (22), JSH-2 (23), KRM-13 (24); B) 4th millennium BP (violet): SNR-4bis/2 (1), SNR-1D (2), OBS-1 (3); C) 3rd millennium BP (red): SNR-3 (1), and JSH-10 (2) (drawings by R. Nisbet and P. Biagi)

headlands of solid massive rocks (f.i. at Gadani, Phuari, Daun and Sonari); and 3) the Indus Delta, corresponding in our survey to the coast from Karachi to Aban Shah, south of Thatta in longitude, and Gharo-Makli Hills (Thatta) to Kot Raja Manjera (Jerruck) in latitude (**Fig. 7**).

⁶³ East et al. 2015.

⁶⁴ Lambrick 1986.

⁶⁵ Inam et al. 2007.

⁶⁶ Clift/Giosan 2014; Clift et al. 2014.

⁶⁷ Wells/Coleman 1984; Meadows/Meadows 1999; Shuaib/Tariq Shuaib 1999.




Fig. 7 Distribution map of the radiocarbon-dated sites from mangrove specimens from Bhambor, in the west, to the Makli Hills, in the east. Garo-1 (1), THR-3 (2), Beri (3), JSH-1bis (4), JSH-2 (5), JSH-10 (6), KKT-2 (7), KKT-4 (8), KKT-3 (9), MKL-10 (10), and MKL-1 (11) (drawing by P. Biagi)




Fig. 8 Lake Siranda: Distribution map of the archaeological sites, mainly shell middens, discovered during the 2010-2014 surveys. 8th millennium BP (light blue), 7th millennium BP (blue), 6th millennium BP (green), 5th millennium BP (yellow), white dots represent undated sites, Balakot (BLK-1) (drawing by R. Nisbet)




Fig. 9 Lake Siranda: Plot of the calibrated dates obtained from mangrove gastropods from the shell middens, and the Chalcolithic/Bronze Age site of Balakot (BLK-1) (Las Bela, Balochistan) (scatterplot by T. Fantuzzi)

1) Siranda palaeo-lagoon

At present a detailed sequence in the use of palaeo-mangroves is known from Lake Siranda.⁶⁸ The early morphological history of the basin is still poorly understood.⁶⁹ It is accepted by the authors that it was formerly part of the present Sonmiani Lagoon (Miāni Hor), sharing with it its dominant environmental aspects and that it “functioned as a tidal lagoon in the not-too-distant past” (Fig. 8).⁷⁰ According to the above author Siranda was formerly connected to Sonmiani Lagoon from which it was later separated by sand dune formations, eventually stabilised by vegetation.⁷¹

The 33 radiocarbon dates so far obtained from the area (Fig. 9) show that the first exploitation of mangroves started ca. one century after the radiocarbon result obtained from the Mulri Hills, east of Karachi (MH-15, GrA-63863: 7320±40 BP), and lasted for the entire 7th and 6th millennia BP. The earliest dates, spanning from the last centuries of the 8th and the middle of the 7th millennium BP, come from sites located along the south-eastern side of the basin, some of which yielded chipped stone assemblages with geometric microliths obtained mainly from Gadani reddish-brown flint (Fig. 10). Just after the mid 7th millennium BP the sites spread toward north-east, at the inner sides of the depression. After this period the shell middens seem to


Fig. 10 Lake Siranda: chipped stone tools of Gadani reddish-brown flint from shell midden SRN-29: trapezoidal geometrics (1–3), lunate (4), backed bladelets (5, 10), micro-drill (6), truncation (7) and unretouched bladelets (8, 9) (from Biagi et al. 2013b fig. 6)

⁶⁸ Minchin 1907: 9; Hughes-Buller 1908: 96.

⁶⁹ Biagi et al. 2013b.

⁷⁰ Snead 1966, 60.

⁷¹ Snead 1969, 34.




disappear from the northern part of the palaeo-lagoon, and are found again at its south-western corner.

Quite rapid shift of sites to its southern margin took place between the end of the 6th and the end of the 5th millennium BP, showing the progressive desiccation of the ancient lagoon, which would have lost its connections with the sea probably by the end of the 5th millennium BP, when the sea-level started to decrease.⁷² The more recent shell midden found at Siranda yielded a date of 4315±35 BP (SRN-57, GrA-57533), which is three centuries more recent than that obtained from a *T. palustris* specimen collected from the surface of Balakot, ca. 10 km to the south-east (BLK-1, GrA-55828: 4660±40 BP). The stratified Chalcolithic/Bronze Age mound of Balakot (Fig. 4) developed during the last phases of exploitation of the Siranda palaeo-mangrove, as shown by many *Terebralia palustris* shells retrieved from the Bronze Age occupation layers.⁷³ In effect

Fig. 11 Daun: Plot of the calibrated dates obtained from mangrove gastropods from the shell middens (scatterplot by T. Fantuzzi)

⁷² Gupta 1972: 360.

⁷³ Dales 1974; Meadow 1979: Fig. 6.



they are very common to the Bronze Age Indus period settlement, where they represent 67% of the total shell assemblage, while their presence reaches only 18% in the Chalcolithic layers.

2) The Balochistan coast between Cape Gadani and the Hab River mouth

South of Sonmiani Lagoon the coast forms a long, low sandy strip, interrupted only by two rocky headlands at Gadani and Phuari. The first belongs to the Bela Ophiolite, a sequence of basaltic pillow-lavas, inter-flow sedimentary rocks (chert, argillite and limestone) mostly of Upper Cretaceous age⁷⁴ emerging as the western part of Mor Range, and the Parh limestone formation (Upper Cretaceous), a unit of the Pab Range.

Ras Gadani is separated from Ras Phuari, some 3.5 km to the south, by a sand strip that runs in a NNE-SSW direction, in the centre of which is a small, seasonal stream called Kunari Dhora, which flows into the Arabian Sea. The Bela Ophiolites outcrop runs along the coast, south of the mouth of the Kunari Dhora as far as Ras Phuari. Two almost identical dates from Gadani (GDN-0, GrN-26369: 4460±30 BP) and Ras Phuari (PHR-11, GrA-55826: 4415±40 BP) prove the existence of mangroves probably at the mouth of the mentioned small streams, around the middle of the 5th millennium BP, a time when Siranda palaeo-lagoon had already transformed into a saline depression, fed only by monsoon rains and seasonal rivers.⁷⁵

Further south, the shell middens of Daun Bay lie partly along the sand beach around and south of the bay, some 4-10 m above the maximum level reached by the tide, partly on the top of the Pleistocene marine terrace (16-20 m) extending south of a small headland.⁷⁶ Their distance from the present shoreline varies from 60 to 700 m. Most sites consist of heaps or scatters of fragmented *Terebralia palustris* gastropods,⁷⁷ although other mangrove and marine species are represented, among which are *Telescopium telescopium* and *Anadara uropygmelana*.

The radiocarbon results from the Daun sites indicate that the exploitation of the mangrove resources was not "continuous" (Fig. 11). It took place mainly during two distinct periods of the middle of the 7th and the 5th millennium BP respectively,⁷⁸ thus pointing to the existence of coastal forests at the same time as at Siranda. The first cluster of Daun dates, belonging to the Neolithic, shows a

⁷⁴ Blanford 1880; Vredenburg 1909; Sarwar 1992.

⁷⁵ Minchin 1907; Biagi et al. 2013b.

⁷⁶ Snead 1966, 47; 1967; 1969, 38; Snead/Frushman 1968, 1673.

⁷⁷ Biagi 2004; Biagi/Franco 2008; Biagi et al. 2012.

⁷⁸ Biagi et al. 2012.

$\delta^{13}\text{C}$ ratio ranging from -3.44 to -3.97, which is compatible with a mixed marine mangrove ecosystem. A more recent sporadic episode of exploitation, which took place during the Chalcolithic (Daun-6, GrN-28802: 5370 \pm 35 BP and Daun-116: GrA-66637: 5360 \pm 40 BP), shows in one case a dramatic increase to +1.27 (GrN-28802), possibly an indicator of environmental stress. The samples obtained from the second main cycle of exploitation of Daun Bay, as well as Capes Gadani and Phuari, show once again lower $\delta^{13}\text{C}$ (-4.49 to -6.10) that are typical of a healthy mangrove ecosystem with mixed marine and freshwater. However, a much more recent result obtained from one single *T. palustris* sample shows that mangroves were still growing immediately south of the bay in historical times (Daun-117, GrN-31494: 1440 \pm 30 BP).

At the southern mouth of the Hab River the southernmost extension of Pab Range rises with its Jhill limestone unit, a member of the Miocene Gaj Formation.⁷⁹ On the top of a saddle 30-40 m high located near the village of Sonari, ca. 7 km north-east of Ras Mauri (Cape Monze) (Fig. 12), local pre-historic fishermen living in rectangular stone structures (Fig. 13) collected mangrove Gastropods in a tidal forest certainly growing along the estuary of the Hab River (Fig. 14), at least since the early 4th millennium BP.⁸⁰ Similar dates were obtained from the Bronze Age Indus Civilisation small settlement of Pir Shah Jurio (PSH-1, GrN-26370: 4130 \pm 20 BP; PSH-1bis, GrA-66638: 4270 \pm 35 BP) located on a protruding terrace along the left, eastern bank of the Hab River, surrounded by alluvium, some 6.5 km north-east of its mouth⁸¹ (Fig. 15). The last two results from Sonari (SNR-3, GrA-62249: 2190 \pm 30 BP and SNR-2, GrA-59834: 670 \pm 50 BP), and one from Daun (Daun-117, GrN-31494: 1440 \pm 30 BP) are so far the only historical dates available for the Pakistani palaeo-mangroves (Fig. 16).




Fig. 12 Sonari: Distribution map of the 5th millennium BP sites radiocarbon-dated from mangrove gastropods: PSH is Pir Shah Jurio (top); distribution map of the 5th millennium BP (yellow), 4th millennium BP (violet), 3rd millennium BP (red), and Historical period (ochre). The white dot is a group of Islamic tombs (bottom) (drawings by P. Biagi)

⁷⁹ Blanford 1880; Naseem et al. 1996.

⁸⁰ Biagi/Nisbet 2014.

⁸¹ Fairservis 1993 Fig. 9.1.



Fig. 13 Pir Shah Jurio: the mature Indus Civilisation site (top), and *Terebralia palustris* specimens from the surface of the same site (bottom) (photographs by P. Biagi 2014)



Fig. 14 Sonari: location of the radio-carbon-dated Bronze Age fishermen site SNR-1, in the centre of the saddle (top), and rectangular stone structures filled with marine and mangrove shells from the same site (bottom) (photographs by P. Biagi 2013)



Fig. 15 Sonari: The Hab River mouth from the south, with the location of the Islamic cemetery (ochre dot) (photograph by P. Biagi 2013)




Fig. 16 Sonari: Plot of the calibrated dates obtained from mangrove gastropods (scatterplot by T. Fantuzzi)

3) The Indus Delta

The date from the Mulri Hills, at the eastern outskirts of Karachi (MH-15, GrA-63863: 7320±40 BP), is the oldest radiocarbon result available to date from mangrove gastropods showing the presence of mangroves along the seaside of present-day Pakistan. The Mulri Hills are a unique location in the entire study region. Their surface was literally covered with Late (Upper) Palaeolithic and Mesolithic sites until the 1980s. The hills are rich in freshwater that gushes out of roughly east-west oriented faults, from which originate small streams that flow southward down into the Malir River and soon after the Ghizri Creek west of Rehri, and Kadiro Creek.⁸² Given the inland position of the hills ca. 70 m high⁸³ it is probable that the *T. palustris* shells were collected from mangroves that flourished along the coast ca. 8 km to the south, where they still survive (Manora Island and surrounding areas) or along the mouth of the Malir River at Ghizri Creek and further south, or east at Kadiro Creek⁸⁴ (Fig. 17). It is important to point out that a date similar to the oldest so far available from the Mulri Hills (MH-15, GrA-63863: 7320±40 BP) comes from a *T. palustris* scatter found at Rehri, a location facing the latter aforementioned creek (RHR-3bis, GrA-66631: 7045±40 BP) (Fig. 18). However, according to Professor A. R. Khan the

whole coastal area around Karachi has been subjected to at least three phases of uplift during the Holocene, with the formation of a series of raised beaches and marine terraces (at 6-7 m, 9-12 m and over 15 m respectively).⁸⁵ How effective these tectonic processes were instrumental in causing changes to the coastal mangroves is still to be understood, as they resulted in the seaward advance of the shoreline.

Moving eastward, the first AMS-dated occurrence of *Telescopium telescopium* is known from Gharo (Garo-1, GrA-59844: 6320±60 BP). Located on a limestone terrace, at an altitude of ca. 30 m, the site is 4-5 km from the ruins of the 8th-13th century AD Gharo Creek outpost of Bhambor,⁸⁶ at present along the shore of an active channel of the Indus River and close to the actual mangrove.

A consistent number of radiocarbon dates has been obtained west and south of Thatta, one of the ancient capitals of Sindh. The shell middens are located some 30 km from the present western coastline, and more than 80 km from the mouth of the main Indus channel, in the south. Almost all the samples come from the top or, less frequently, the side of isolated tracts of calcareous and sandstone hillocks, rising from the alluvial plain between 10 and 30 m, at an elevation of 15-40 m above the sea level. These features were undoubtedly surrounded by the sea before the advance of the Delta, forming

⁸² Khan 1979b; Biagi 2003-2004, 2017.


⁸³ Zaidi et al. 1999.

⁸⁴ Snedaker 1984; Kazi 1999.

⁸⁵ Khan 1979a, 19-21.

⁸⁶ Majumdar 1934, 19; Cousens 1998, 64; Felici et al. 2016.

Fig. 17 Mulri Hills: Distribution map of the sites at the eastern outskirts of Karachi. The other black dots show the location of some archaeological sites discovered by Professor A. R. Khan in the late 1970s (drawing by P. Biagi)



an archipelago which could be easily reached from the coast even in historical times. Arrian's account on Nearchos journey mentions some "isles" when the fleet reached the sea. The more evident of these elevations are the Makli Hills, running north-south with the ruins of an impressive fortress known as Kalan Kot (8th to 18th century according to Cousens),⁸⁷ close to which the earliest date for the area has been obtained (KKT-2, GrN-32464: 6320±45 BP) (Fig. 19-21).

The results yielded by nine sites show the presence and exploitation of mangroves since the mid 8th to the end of the 6th millennium BP. There are, however, interesting exceptions from Shah Husein (JSH-1bis, GrA-66636: 5800±40 BP; JSH-2, GrA-45181: 4245±40 BP and JSH-10, GrA-62255: 2715±30 BP), an isolated rocky cliff ca. 13 km west-southwest from Thatta, from which we have evidence of several mangrove shell scatters as well as chipped stone artefacts (Fig. 22). The latter date shows that a mangrove environment lasted locally probably as late as the Hellenistic period, along one or more creeks connecting the site to the seaside over a period of four millennia.

Some 13 km west of Thatta, and 3 south-west of Gujo, another limestone terrace is well known in the archaeological literature as a fortified Amri Culture site, from the surface of which thousands of chipped stone tools have been retrieved.⁸⁸ The site, known as Tharro Hills, is located at the south-eastern edge of the terrace. It is surrounded by two parallel, semi-circular stone walls (Fig. 23). During the intensive survey carried out in January 2008,⁸⁹ many specialised areas were recorded, 41 of which yielded characteristic Amri chipped stone tools, among which are bladelets with semi-abrupt retouch, truncations and typical elongated scalene triangles⁹⁰ (Fig. 24), and a few characteristic painted potshers. Two radiocarbon dates were obtained from samples of *Ostreidae* (THR-1, GrN-27053: 5240±40 BP) and *T. palustris* (THR-3, GrA-47084: 5555±35 BP) respectively recovered from a well-defined spot of shells located along the southern edge of the inner wall, confirming once again attribution of the site to the Chalcolithic.

South of the Tharro Hills another Chalcolithic site was discovered at the top of a small limestone terrace rising from the Indus alluvium, locally called Beri.⁹¹ A *Terebralia palustris* shell sample was collected from the surface of this site, later radiocarbon-dated to 5960±50 BP (Beri-1, GrN-32166)




Fig. 18 Indus Delta: Plot of the calibrated dates obtained from mangrove gastropods from the shell middens (scatterplot by T. Fantuzzi)

The only dated site on the left bank of Indus is an isolated rocky hill known as Aban Shah (Fig. 25). The site lies some 70 km north of the present Arabian Sea shore. The date (ABS-1, GrA-47082: 3790±35 BP) provides evidence of a local mangrove still growing after the end of the Bronze Age. It shows that 1) the progress of the coastline has been relatively slow (20 km/3500 years) in comparison with other points of the Delta; 2) the presence of mangroves surrounding the site during the early 4th millennium BP would exclude it as the island "out in the sea" of the 4th century BC reported by Arrian, erroneously considered by H. T. Lambrick⁹² the islet on which Alexander landed, before sailing back to Pattala.⁹³

⁸⁷ Cousens 1998.

⁸⁸ Majumdar 1934; Piggott 1950; Khan 1979a, 5; Biagi 2005.

⁸⁹ Biagi/Franco 2008.

⁹⁰ Biagi 2005.

⁹¹ Biagi 2010, 9.

⁹² Lambrick 1986, 113.

⁹³ see Eggermont 1975 Map 2.



Fig. 19 Makli Hills: The site of Kalan Kot 2 (KKT-2) (top), and the eastern edge of the hills, along the central part of the terrace looking at the Indus alluvial plain (bottom) (photographs by P. Biagi 2011)



Fig. 20 Makli Hills: The site of Kalan Kot 4 (KKT-4) (top), and fragments of *Telescopium telescopium* mangrove gastropods on its surface (bottom) (photographs by P. Biagi 2012)



Fig. 21 Makli Hills: The site of MKL-10 (blue spot) (top), and fragments of marine shells on its surface (bottom) (photographs by P. Biagi 2013)



Fig. 22 Shah Husein: The hill from north-west (top), and the radiocarbon-dated site of JSH-1bis from which one trapezoidal arrowhead of *fleche trenchant* type has been recovered (bottom) (photographs by P. Biagi 2009)




Fig. 23 Tharro Hills: Location of the fortified Amri Culture site (red square), the radiocarbon-dated Chalcolithic (THR-3: green dot) and Neolithic (THR-2: blue dot) sites (top), and the fortified Amri Culture site from the east (bottom) (from Biagi and Franco 2008 Fig. 7, with modifications; photograph by P. Biagi 2009)




Fig. 24 Tharro Hills: characteristic long triangles of the Chalcolithic Amri culture (from Biagi 2005 Fig. 7 with modifications)



Fig. 25 Aban Shah: A view of some hillocks from the highest limestone terrace (photograph by P. Biagi 2011)

In this chronological frame, the most intriguing date obtained from our survey in the Delta area comes from an inland site located on a flat-topped limestone mesa (45-47 m asl) of the Kirthar formation, called Kot Raja Manjera. The site is famous for its Buddhist stupa and other archaeological remains attributed to the 5th century AD (**Fig. 26**). The terrace is roughly east-west oriented, along the south-western bank of an ancient meander of the Indus, which at present flows some 5 km to the east. Kot Raja Manjera is a fortified Amri Culture Chalcolithic settlement. From its surface also a few Bronze Age potsherds were collected as well as many chipped stone tools among which are small drills for bead manufacture.⁹⁴ A few marine shells were recovered as well as one *T. palustris* specimen, which was AMS-dated to 4635±35 BP (KRM-13, GrA-47083). Close to the terrace the river forms a semicircular bend that in prehistoric times lapped the limestone formation on which the village of Lakho Pir is located.⁹⁵

According to the above results we have to admit that 1,000-1,500 years after the exploitation of the mangroves in Thatta region the same was still happening around a site ca. 60 km northward. In fact Kot Raja Manjera yielded the northernmost finding of a mangrove shell within the entire Indus Delta region.

Calibration problems

The problems related with the calibration of radiocarbon results of both mangrove and marine specimens from shell middens excavated along the shores of the Oman Peninsula Arabian Sea coasts have already been discussed in several papers with contrasting results.⁹⁶ In the case of Las Bela and Sindh, the nearest data as to deep-water upwelling and/or oceanic reservoir that might affect the calibration of the radiocarbon dates from the study area come from a sea core off the Makran coast⁹⁷ some 300 km north-west of Port Okha in Gujarat.⁹⁸ This value, chosen to calibrate the dates presented in **Table 1**, should be treated with caution, as both local and diachronic variations in the ΔR that have proven to be relevant in modern age samples⁹⁹ might be unrecognisable on the basis of the available data.

Moreover, as the $\delta^{13}\text{C}$ values show (**Table 1**), the samples presented in this paper come from mangrove environments, from which we cannot exclude the presence of old carbon caused by variations of the percentage of fresh vs. marine water, hard water effects, and organic litter composition. However, their percentage is very low off the Las Bela coast.¹⁰⁰

⁹⁶ Uerpman 1991; Biagi 1994; Saliège et al. 2005; Zazzo et al. 2012; 2016.

⁹⁷ 56KA: von Rad et al. 1999; Saliège et al. 2005 Fig. 1.

⁹⁸ 229±27 ¹⁴C years: Reimer/Reimer 2001.

⁹⁹ Dutta et al. 2001.

¹⁰⁰ Stewart/Pilkey 1966 Fig. 9.

⁹⁴ Khan 1979c, 72.

⁹⁵ Biagi 2010 Fig. 14.




Fig. 26 Kot Raja Manjera: Site location (red square) and point 13 (yellow dot) from which one *Terebralia palustris* fragment was radiocarbon-dated (GrA-47083) (top). The other black dots show concentrations of Chalcolithic and Bronze Age finds (bottom) (drawing by P. Biagi and R. Nisbet)

Discussion

Leaving apart the 9th millennium BP date obtained from the Kadeji River marine bivalve discussed below (KDJ-1), according to the aforementioned results, around the end of the 8th millennium BP there is evidence of mangroves at or near the mouth of a few rivers in the Sonmiani and Karachi areas, which were exploited by the earliest Holocene inhabitants of the northern coast of the Arabian Sea. These data can be compared with those obtained from the earliest shell middens of the coasts of Oman,¹⁰¹ and more generally the entire coastline of the Arabian Peninsula, from which we have a reasonable set of dates confirming that the territory began to be settled roughly during this period.¹⁰² According to the available radiocarbon results from mangrove shells, between the 7th and 6th millennia BP the coast of the Indus Delta was located along an arc running from Manora-Ghizri Creek, in the west, to Thatta-Makli Hills, in the east. It seems, therefore, that the western sector of the Delta, from Karachi to Bhambor-Gharo developed at that time, and later seaward accretion occurred in a minor extension.¹⁰³ In contrast, the central part of the Delta, south of Thatta, has been subjected to a much larger advance even in historic times, though it is not possible, on the basis of our data, to establish the dynamics in the course and position of the palaeo-channels.¹⁰⁴

With regard to the last two millennia, since the 18th century AD many authors have tried to reconstruct the road followed by Nearchos fleet in its journey back to Mesopotamia, in relationship with the location of the ancient seashore, generally using as a source the itineraries handed down by Greek and Roman historians.¹⁰⁵ According to H.R. Haig¹⁰⁶ the head of the Delta would be situated at the latitude of Thatta in Hellenistic times, which well agrees with the radiocarbon data. A similar opinion is shared by T.H. Lambrick¹⁰⁷ yet suggesting that the western coast of the Delta was not far from the Makli Hills and Pir Patho (Thatta) in Alexander's times. This hypothesis is not consistent with both the radiocarbon dates, and the reconstruction by P. H. L. Eggermont,¹⁰⁸ though his interpretation of Aban Shah hillock as "the island in the sea"¹⁰⁹ is not confirmed by our *T. palustris* date (OBS-1, GrA-47082: 3790±35 BP). In contrast it points to the presence of

mangroves in the area at least one thousand years before Alexander's retreat from India.

According to D.A. Holmes, deltaic morphologies are found as inland as to 55 km north-east of Hyderabad in historical times, "suggesting a very high rate of alluvial aggradation and delta advancement".¹¹⁰ Similarly the reconstruction proposed by L. Flam,¹¹¹ which is mostly based on aerial photography and the distribution of archaeological sites in the Delta, suggests that the 6th-5th millennium BP coast was probably located somewhere between Hyderabad and Thatta,¹¹² an hypothesis that would better fit with our dates, and could also explain such an early date as that from Kot Raja Manjera (KRM-13, GrA-47083: 4635±35 BP).

Conclusion

The results obtained from the 2000-2014 surveys carried out along the coast of Lower Sindh and Las Bela in Balochistan, have shown the great archaeological potential of the area for the study of the prehistory of the two territories. The discovery of an impressive number of sites, and their radiocarbon dating, has shown that the earliest settlements of the coastal strip are to be referred to the last centuries of the 8th millennium BP. Furthermore the data retrieved from our research has shown that

1. Early Neolithic settlements are not exclusively limited to the regions of the interior of Pakistani Balochistan¹¹³ as often suggested.¹¹⁴ According to both the new radiocarbon results, and the analysis of the lithic assemblages recovered from the Siranda shell middens, among which are geometric microliths obtained from Gadani reddish-brown flint (Fig. 10), Neolithic sites undoubtedly existed along the ancient coastline. Our results show that they are more or less contemporary, or slightly more recent, to the earliest occupation of Mehrgarh in the Kachi Plain,¹¹⁵ though the radiocarbon chronology of the aceramic Neolithic occupation of this site is very controversial.¹¹⁶ This fact opens new perspectives to the interpretation of the Neolithic archaeology of Balochistan at present known from a far too small number of sites.¹¹⁷
2. Movements along the northern coast of the Arabian Sea began at least around the beginning

¹⁰¹ Berger et al. 2013; Zazzo et al. 2016.

¹⁰² Boivin/Fuller 2009.

¹⁰³ Tremeneheere 1867.

¹⁰⁴ Wilhelmy 1968.

¹⁰⁵ Eggermont 1975; Kevran 1995; Baynham 2005; Biagi 2017.

¹⁰⁶ Haig 1894.

¹⁰⁷ Lambrick 1986, 113.

¹⁰⁸ Eggermont 1975.

¹⁰⁹ Eggermont 1975 Map. 2.

¹¹⁰ Holmes 1968, 369.

¹¹¹ Flam 1999.

¹¹² Flam 1999, 65.

¹¹³ Fairervis 1956; Jarrige 2004.

¹¹⁴ See also Possehl 2003 Fig. 2.5.

¹¹⁵ for a comparison see Petrie et al. 2010 Table 2.1.

¹¹⁶ Jarrige/Lechevallier 1980; Jarrige et al. 1995, 555; Lechevallier 2003, 15, note 4.

¹¹⁷ Fairervis 1956; Shaffer 1978; Petrie et al. 2010.

of the 7th millennium BP. A scatter of oyster shells discovered along the southern edge of the Tharro Hills yielded a date of 6910±60 BP (THR-2, GrN: 32119). It can be compared with some of the most ancient results from Lake Siranda (see Fig. 9: SRN-56, GrA-57702: 6980±35 BP and SRN-33, GrA-54291: 6770±35 BP respectively), while the Makli Hills were first settled during the second half of the same millennium (KKT-2, GrN-32464: 6320±45 BP, and MKL-10, GrA-62256: 6140±40 BP) (Fig. 20 and 21). This would suggest that navigation along the northern coast of the Arabian Sea had already started in this period, if not already a few centuries before,

3. The Indus Delta “islands” were undoubtedly exploited since the Late Neolithic and Chalcolithic periods as shown by the presence of Amri Culture sites, like the Tharro Hills, and the radiocarbon results from Beri and the Makli Hills (Fig. 18). The same sites do not seem to have been settled during the Bronze Age Mature Indus Civilization. This fact is intriguing, since we know that during the mid 3rd millennium BC, Indus sailors and traders systematically moved across the Arabian Sea to reach the coasts of the Arabian Peninsula, where Indus outposts are known since a few decades,¹¹⁸

4. The series of radiocarbon dates obtained from shells provide excellent arguments for a preliminary reconstruction of the Holocene history of the mangroves along the coasts of Las Bela, and the variation of the profile of the Indus Delta during the same period. In this region mangroves followed the millennial built-up of the land at least since the 6th millennium BP. Unfortunately at present no dates are available for the earliest formation of the Delta, undoubtedly pre-Holocene,¹¹⁹ whose apex several authors would set dozens km north to Hyderabad. The advance of the Delta coast during the last millennium could by no means hinder the use of the river for sailing northward, though the number of 5,000 boats moving from the ancient port of Debal (al-Daybul, most probably Bhambor¹²⁰) at the time of Sultan Feroz Shah Tughluq invasion of Sindh (1365-1367 AD)¹²¹ might be exaggerated by the ancient geographers. According H. G. Raverty, near Thatta the river was so large that from a side of the river “the land around could not be distinguished”¹²² and therefore “a great part of the delta south of [Thatta] has been formed since these events happened”.¹²³

5. The new radiocarbon results (Table 1) show that mangroves flourished during the mid-Holocene along the northern coast of the Arabian Sea. For

still unknown reasons they were no longer exploited after the 5th millennium BP at Siranda, or much later, around the end of 3rd millennium BP at Sonari, and very rarely even in historical times (Daun and Sonari). At the present state of the research it is impossible to say whether this fact is related to their disappearance, as a consequence of eco-climatic changes, as should be the case for Lake Siranda, or cultural reasons. The data achieved from fieldwork show that all the palaeo-mangroves so far recorded at Las Bela (Siranda, Gadani, Phuari and Daun), as well as those still flourishing one or two centuries ago at the mouth of the Hab River (SNR-2, GrA-59834: 670±50 BP) have now totally disappeared. However, most of the radiocarbon dates fall into a period comprised between two dramatic climatic changes. The first took place during the second half of the 9th millennium BP, following a sudden reduction of precipitations all over this part of the Indian Subcontinent,¹²⁴ while the second, a more drastic drought, occurred around 4200 BP,¹²⁵ and most probably contributed to the decline of the Indus Civilization.¹²⁶ This last climatic crisis seems well represented in the sharp fall of occurrences in the distribution map between the 4th and 3rd millennium BP (Fig. 8 and 9).

6. A new radiocarbon result from a site re-discovered along the southern bank of the Kadeji River at its confluence with the Mol, suggests that mangroves were already present somewhere along the coast already during the second half of the 9th millennium BP (KDJ-1, GrA-63862: 8275±45 BP) (Fig. 5A-5). Though the assay was obtained from a fragment of marine bivalve, its negative $\delta^{13}\text{C}$ value of -4,44, perfectly fits into the general picture of a mangrove environment.¹²⁷ This fact would indicate that already during the Mesolithic period, the area surrounding the Malir River, was partly covered with mangroves, which were exploited by groups of last hunter-gatherers.

Acknowledgements

The research in Lower Sindh and Las Bela was made possible thanks to the financial support of the Italian Ministry of Foreign Affairs (MAE, Rome), the Archaeology Research Funds of Ca' Foscari University (Venice), the EURAL Gnutti spa (Rovato, Brescia), and the logistic support of Sindh University, Jamshoro, and the University of Balochistan, Quetta. Special thanks are due Mr. A.

¹¹⁸ Charpentier 1996; Méry 1996; 2000; Ray 2003; Ratnagar 2004.

¹¹⁹ Clift et al. 2014, 185

¹²⁰ Pathan 1978, 417.

¹²¹ Panhwar 1983, 32.

¹²² Raverty 1895, 126.

¹²³ Raverty 1895, 126, note 235.

¹²⁴ Staubwasser et al. 2002.

¹²⁵ Staubwasser et al. 2003; Dixit et al. 2014.

¹²⁶ Gangal et al. 2010.

¹²⁷ Biagi 2018

Girod (Italian Malacological Society) who took part in the 2012-2013 fieldwork seasons in Las Bela, Prof. Mazharul Haq Siddiqui (former Vice-Chancellor of Sindh University, Jamshoro), Mr. Shoukat Shoro (former Director of the Institute of Sindhology, Jamshoro), Mir Atta Mohammad Talpur, Mir Ahmed

Farooq Talpur, Mir Ghulam Rasool Talpur, and Mir Abdul Rehman Talpur for all their help, support and friendship. Special thanks are due to Prof. K. Thomas (University College, London) for the revision of the English text, the useful comments and suggestions.

Bibliography

- Aitchison 1869
J. E. T. Aitchison, A Catalogue of the Plants of the Punjab and Sindh (London 1869) (reprinted by Bishen Singh Mahendra Pal Singh, Dehra Dun, India, 1982).
- Aitken 1907
E. H. Aitken, Gazetteer of the Province of Sind (Karachi 1907).
- Ambaglio 1994
D. Ambaglio (ed.), *Anabasi di Alessandro* (Milano 1994).
- Amir Khanov et al. 2001
Kh. A. Amir Khanov/B. Vogt/A. V. Sedov/V. Buffa, Excavation of a settlement of prehistoric fishermen and mollusk gatherers in the Khor Umayra lagoon, Gulf of Aden, Republic of Yemen. *Archaeology Ethnology and Anthropology of Eurasia* 4:8, 2001, 2–12.
- Amjad/Kamaruzaman 2007
A. S. Amjad/J. Kamaruzaman, Mangrove Conservation through Community Participation in Pakistan: The Case of Sonmiani Bay. *International Journal of Systems Applications, Engineering and Development* 1:4; 2007, 75–81.
- Amjad et al. 2007
A. S. Amjad/I. Kasawani/J. Kamaruzaman. Degradation of Indus Delta Mangroves in Pakistan. *International Journal of Geology* 3:1, 2007, 27–34.
- Baig/Iftikhar 2006
S. P. Baig/U. A. Iftikhar, Are the Mangroves for the Future? Empirical evidence of the value of Miani Hor Mangrove Ecosystem as the basis for investments. Pakistan, IUCN, 2006.
- Bailey/Parkington 1988
G. Bailey/F. Parkington, The archaeology of prehistoric coastlines: an introduction. In: G. Bailey/F. Parkington (eds.), *The Archaeology of Prehistoric Coastlines* (Cambridge 1988), 1–10.
- Bailey et al. 2013
G. N. Bailey/M. G. Meredith-Williams/A. A. Alsharek, Shell Mounds of the Farasan Islands, Saudi Arabia. In: G. N. Bailey/K. Hardy/A. Camara (eds.), *Shell Energy. Mollusc Shells as Coastal Resources* (Oxford-Oakville 2013) 241–254.
- Baillie 1890
A. F. Baillie, *Kurrachee Past, Present and Future* (Calcutta 1890).
- Bar-Yosef Mayer/Beyn 2009
D. E. Bar-Yosef Mayer/A. Beyn, Late Stone Age Shell Middens on the Red Sea Coast of Eritrea. *Journal of Island and Coastal Archaeology* 4:1; 2009, 108–124.
- Bar-Yosef Mayer/Beyn 2013
D. E. Bar-Yosef Mayer/A. Beyn, Mid-Holocene Shell Middens in Eritrea. In: G. N. Bailey/K. Hardy/A. Camara (eds.), *Shell Energy. Mollusc Shells as Coastal Resources* (Oxford-Oakville 2013) 229–239.
- Baynham 2005
E. Baynham, Arrian's sources and reliability. In: J. Romm (ed.), *The Landmark Arrian. The Campaigns of Alexander* (New York 2005) 325–332.
- Berger et al. 2013
J. F. Berger/V. Charpentier/R. Crassard/C. Martin/G. Davtian/J. A. López-Sáez, The dynamics of mangrove ecosystems, changes in sea level and the strategies of Neolithic settlements along the coast of Oman (6000–3000 cal. BC). *Journal of Archaeological Science* 40, 2013, 3087–3104.
- Berger et al. 2005
J.-F. Berger/S. Cleuziou/G. Davtian/M. Cattani/F. Cavulli/V. Charpentier/M. Cremaschi/J. Giraud/P. Marquis/C. Martin/S. Méry/J.-C. Plaziat/J.-F. Saliège, *Évolution Paléogéographique du Ja'alan (Oman) à l'Holocène Moyen: Impact sur l'évolution des Paléomilieux Littoraux et les Stratégies d'Adaptation des Communautés Humaines*. *Paléorient* 31, 2005, fasc.1, 46–63.
- Besenal 1992
R. Besenal, Recent Archaeological Surveys in Pakistani Makran. In: C. Jarrige (ed.), *South Asian Archaeology 1989 Monographs in World Archaeology* 14 (Madison 1992) 25–35.
- Biagi 1988
P. Biagi, Surveys along the Oman Coast: Preliminary Report on the 1985–1988 Campaigns. *East and West* 38, 1988, fasc. 1–4, 271–291.
- Biagi 1994
P. Biagi, A radiocarbon chronology for the aceramic shell-middens of coastal Oman. *Arabian Archaeology and Epigraphy* 5, 1994, 17–31.
- Biagi 2003–2004
P. Biagi, The Mesolithic Settlement of Sindh: A Preliminary Assessment. *Praehistoria* 4–5, 2003–2004, 195–220.
- Biagi 2004
P. Biagi, New radiocarbon dates for the prehistory of the Arabian Sea coasts of Lower Sindh and Las Bela in Balochistan. *Rivista di Archeologia* 28, 2004, 5–16.
- Biagi 2005
P. Biagi, The chipped stone assemblage of the Tharro Hills (Thatta, Sindh, Pakistan): a preliminary typological analysis. In: F. Martini (ed.), *Askategi miscellanea in Memoria di Georges Laplace*. *Rivista di Scienze Preistoriche Supplemento* 1, 2005, 553–566.
- Biagi 2008
P. Biagi, The shell-middens of the Arabian Sea and Gulf: maritime connections in the seventh millennium BP? In: A. R. Al-Ansary/K. I. Al-Muaikel/A. M. Alsharek (eds.),

- The City in the Arab World in Light of Archaeological Discoveries: Evolution and Development (Riyadh 2008) 7–16.
- Biagi 2010
P. Biagi, Archaeological Surveys in Lower Sindh: Preliminary Results of the 2009 Season. *Journal of Asian Civilizations* 33:1, 2010, 1, 1–42.
- Biagi 2011
P. Biagi, Changing the prehistory of Sindh and Las Bela coast: twenty-five years of Italian contribution. *World Archaeology* 43:4, 2011, 527–537, 2011, 523–537.
- Biagi 2013
P. Biagi, The shell middens of Las Bela coast and the Indus delta (Arabian Sea, Pakistan). *Arabian Archaeology and Epigraphy* 24, 2013, 9–14.
- Biagi 2017
P. Biagi. *Uneasy Riders: With Alexander and Nearchus from Pattala to Rhambakia*. In: C. Antonetti/P. Biagi (eds.), *With Alexander in India and Central Asia. Moving East and Back to West* (Oxford 2017) 255–278.
- Biagi 2018
P. Biagi, The Mesolithic settlement of Sindh (Pakistan): New evidence from the Khadeji River course. *Præhistoria* 11–12, 2018, 59–74.
- Biagi/Franco 2008
P. Biagi/C. Franco, Ricerche Archeologiche in Balochistan e nel Sindh Meridionale (Pakistan). In: S. Gelichi (ed.), *Missioni archeologiche e progetti di ricerca e scavo dell'Università Ca' Foscari-Venezia, VI Giornata di Studio* (Rome 2008) 9–18.
- Biagi/Nisbet 1992
P. Biagi/R. Nisbet, Environmental history and plant exploitation at the aceramic sites of RH5 and RH6 near the mangrove swamp of Qurum (Muscat – Oman). *Bulletin de la Société Botanique Française* 139:2–4, 1992, 571–578.
- Biagi/Nisbet 2006
P. Biagi/R. Nisbet, The prehistoric fisher-gatherers of the western coast of the Arabian Sea: a case of seasonal sedentarization? *World Archaeology* 38:2, 2006, 220–238 2006.
- Biagi/Nisbet 2014
P. Biagi/R. Nisbet, Sonari: A Bronze Age fisher-gatherer settlement at the Hab River mouth (Sindh, Pakistan). *Antiquity Project Gallery* 341, September 2014. <http://antiquity.ac.uk/projgall/biagi341>.
- Biagi et al. 2012
P. Biagi/T. Fantuzzi/C. Franco, The shell middens of the Bay of Daun: environmental changes and human impact along the coast of Las Bela (Balochistan, Pakistan) between the 8th and the 5th millennium BP. *Eurasian Prehistory* 1:1–2, 2012, 29–49.
- Biagi et al. 2013a
P. Biagi/R. Nisbet/A. Girod, The Archaeological Sites of Gadani and Phuari Headlands (Las Bela, Balochistan, Pakistan). *Journal of Indian Ocean Archaeology* 9, 2013, 75–86.
- Biagi et al. 2013b
P. Biagi/R. Nisbet/A. Girod/T. Fantuzzi, The Middle Holocene mangrove shellfish gatherers of Las Bela coast (Balochistan, Pakistan): new AMS dates from Lake Siranda shell middens. *Antiquity Project Gallery* 087, 2013, 337.
- Biagi et al. 2013c
P. Biagi/A. Girod/R. Nisbet, Prehistoric shell middens, seascapes and landscapes at Lake Siranda (Las Bela, Balochistan). Preliminary results of the 2011 fieldwork season. *Journal of Asian Civilizations* 36, 2013, fasc. 1, 1–25.
- Bjerck 2017
H. B. Bjerck, Settlements and Seafaring: Reflections on the Integration of Boats and Settlements Among Marine Foragers in Early Mesolithic Norway and the Y'amana of Tierra del Fuego. *Journal of Island and Coastal Archaeology* 12:2, 2017, 276–299.
- Blanford 1880
W. T. Blanford, *The Geology of Western Sind*. *Memoirs of the Geological Survey of India* 17, (Calcutta 1880) 1–211.
- Blatter et al., 1929
E. Blatter/C. McCann/T. S. Sabnis, *The flora of the Indus delta* (Madras 1929).
- Boivin/Fuller 2009
N. Boivin/D. Fuller, Shell Middens, Ships and Seeds: Exploring Coastal Subsistence, Maritime Trade and the Dispersal of Domesticates in and Around the Ancient Arabian Peninsula. *Journal of World Prehistory*, 22, 2009, 113–180.
- Campbell 1999
A. C. Campbell, *The Mangrove Communities of Karachi, Present States and Future Prospects*. In: A. Meadows/P. Meadows (eds.) *The Indus River. Biodiversity, Resources, Humankind* (Karachi 1999) 31–41.
- Carter/Crawford 2010
R. Carter/H. Crawford (eds.), *Maritime Interactions in the Arabian Neolithic. Evidence from H3, As-Sabiyah, an Ubaid-related site in Kuwait* (Leiden 2010).
- Cattani/Bökönyi 2002
M. Cattani/S. Bökönyi, *Ash-Shumah. An Early Holocene Settlement of Desert Hunters and Mangrove Foragers in teh Yemeni Tihama*. In: S. Cleuziou/M. Tosi/J. Zarins (eds.), *Essays on the Late Prehistory of the Arabian Peninsula*. *Serie Orientale Roma, XCIII*, 2002, 31–54.
- Charpentier 1996
V. Charpentier, *Archaeology of the Erythraean Sea: Craft Specialization and Resources Optimization as Part of the Coastal Economy on Eastern Coastlands of Oman During the 4th and 3rd Millennia BC*. In *Colloquium XXXII. Trade as a Subsistence Strategy. Post Pleistocene Adaptations in Arabia and Early Maritime Trade in the Indian Ocean*. XII International Congress of Prehistoric and Protohistoric Sciences. *Forlì 1996, A.B.A.C.O.*, 181–192.
- Choquet 1980
M. Choquet, *Determination malacologique de la faune des mollusques marins du Qatar*. *Mission Archéologique Française à Qatar. Tome 1* (Doha 1980) 33–49.
- Cleuziou 2004
S. Cleuziou, *Pourquoi si tard? Nous avons pris un autre chemin. L'Arabie des chasseurs-cueilleurs de l'Holocène au début de l'Age du Bronze*. In: J. Guilaine, (ed.), *Aux marges des grands foyers du Néolithique. Péripéries débitrices ou créatrices?* (Paris 2004) 123–148.
- Clift/Giosan 2014
P. D. Clift/L. Giosan, *Sediment fluxes and buffering in the post-glacial Indus basin*. *Basin Research* 26, 2014, 369–386.

- Clift et al. 2014
P. D. Clift/ L. Giosan/T. J. Henstock/A. R. Tabrez, Sediment storage and reworking on the shelf and in the Canyon of the Indus River-Fan System since the last glacial maximum. *Basin Research* 26, 2014, 183–202.
- Coningham/Young 2015
R. Coningham/R. Young, *The Archaeology of South Asia. From the Indus to Asoka, c. 6500 BCE-200 CE* (New York 2015).
- Cousens 1998
H. Cousens, *The Antiquities of Sind, with Historical Outline* (Karachi 1998) (3rd edition).
- Dales 1974
G. F. Dales, Excavations at Balakot, Pakistan. *Journal of Field Archaeology* 1, 1974, fasc. 1-2, 3–22.
- Dales 1979
G. F. Dales, The Balakot Project: Summary of Four Years Excavations in Pakistan. In: M. Taddei (ed.), *South Asian Archaeology* (Naples 1977) 241–274.
- Dales 1982
G. F. Dales, Adaptation and Exploitation at Harappan Coastal Settlements. In: S. Pastner/L. Flam. (eds.), *Anthropology in Pakistan: Recent Socio-Cultural and Archaeological Perspectives* (Karachi 1982) 154–165.
- Dales/Lipo 1992
G. F. Dales/C. P. Lipo, Explorations on the Makran Coast, Pakistan. *A Search for Paradise. Contributions of the Archaeological Research Facility* 50 (Berkeley 1992).
- Damhoureyeh/Ghalib 2014
S. A. Damhoureyeh/S. A. Ghalib, An overview of the status and distribution of the mangrove forests and their wildlife in Sindh. *Canadian journal of pure and applied sciences* 8:3, 2014, 3051–3055.
- Dixit et al. 2014
Y. Dixit/D. A. Hodell/C. A. Petrie, Abrupt weakening of the summer monsoon in northwest India ~4100 yr ago. *Geology* 42:4, 2014, 339–342. doi: <http://dx.doi.org/10.1130/G35236.1>.
- Dutta et al. 2001
K. Dutta/Bhushan/B. L. K. Somayajulu, ΔR Correction Values for the Northern Indian Ocean. *Radiocarbon* 43, 2001, fasc. 2A, 483–488.
- East et al. 2015
A. Y. East/P. D. Clift/A. Carter/A. Alizai/S. Van Laningham, Fluvial-Eolian Interactions in Sedimentary Signal Buffering: An Example from the Indus Basin and Thar Desert. *Journal of Sedimentary Research* 85, 2015, 715–728.
- Eggermont 1975
P. H. L. Eggermont, Alexander's Campaigns in Sind and Baluchistan and the Siege of the Brahmin Town of Harmatelia. *Orientalia Lovaniensia Analecta* 3 (Leuven 1975).
- Ellison 2014
J. C. Ellison, Vulnerability of Mangroves to Climate Change. In: I. Faridah-Hanum/A. Latiff/K. R. Hakeem/M. Ozturk (eds.), *Mangrove ecosystems of Asia* (New York 2014) 213–231.
- Erlandson/Fitzpatrick 2006
J. M. Erlandson/S. M. Fitzpatrick, Oceans, Islands, and Coasts: Current Perspectives on the Role of the Sea in Human Prehistory. *Journal of Island and Coastal Archaeology* 1:1, 2006, 5–32.
- Erlandson/Rick 2008
J. M. Erlandson/T. C. Rick, Archaeology, Marine Ecology, and Human Impacts on Marine Environments. In: T. C. Rick/J. M. Erlandson (eds.), *Human Impacts on Ancient Marine Ecosystems A Global Perspective* (Berkeley-Los Angeles 2008) 1–19.
- Fairservis 1956
W. A. Fairservis, Jr., Excavations in the Quetta Valley, West Pakistan. *Anthropological Papers of the American Museum of Natural History* 45, 1956, fasc. 2.
- Fairservis 1975
W. A. Fairservis, Jr., *The Roots of Ancient India* (Chicago and London 1975), (2nd Edition).
- Fairservis 1993
W. A. Fairservis, Jr., Allahdino: An Excavation of a Small Harappan Site. In: G. L. Possehl (ed.), *Harappan Civilization. Second Revised Edition* (New Delhi-Bombay-Calcutta 1993) 107–112.
- Felici et al. 2016
A. C. Felici/A. Fusaro/A. Ibrahim/K. Lashari/N. Manasero/M. Piacentini/V. Piacentini Fiorani/A. Tilia, Archaeological excavations at Banbhore, Sindh. Preliminary report of the Pakistani-Italian 2014 and 2015 field seasons. *Parthica* 18, 2016, 125–173.
- Feulner 2006
G. R. Feulner, Occurrence of large mangrove mud creeper *Terebralia palustris* (Linnaeus, 1767) (Gastropoda; Potamodidae) within the Arabian Gulf, and the near Qeshm Island, Iran, in the Strait of Hormuz. *Tribulus* 16, 2006, fasc. 2, 32.
- Flam 1987
L. Flam, Recent Explorations in Sind: Paleogeography, Regional Ecology and Prehistoric Settlement Patterns. *Sindhological Studies*, Sumer 1987, 5–32.
- Flam 1999
L. Flam, The Prehistoric Indus River System and the Indus Civilization in Sindh. *Man and Environment* 24, 1999, fasc. 2, 35–69.
- Gangal et al. 2010
K. Gangal/M. N. Vahia/R. Adhikari, Spatio-temporal analysis of the Indus urbanization. *Current Science* 98, 2010, fasc. 6, 846–852.
- Giosan et al. 2006
L. Giosan/S. Constantinescu/P. D. Clift/A. R. Tabrez/M. Danish/A. Inam, Recent morphodynamics of the Indus delta shore and shelf. *Continental Shelf Research* 26, 2006, 1668–1684.
- Gupta 2004
A. K. Gupta, Monsoon Environments and the Indian Ocean Interaction Sphere in Antiquity: 3000 BC-AD 300. In: Y. Yasuda/V. Shinde (eds.), *Monsoon and Civilization* (Singapore 2004) 133–160.
- Gupta et al. 2003
A. K. D. Gupta/M. Anderson/J.T. Overpeck, Abrupt changes in the Asian southwest monsoon during the Holocene and their links to the North Atlantic Ocean. *Nature* 421, 2003, 354–357.
- Gupta 1972
S. K. Gupta, Chronology of the Raised Beaches and Inland Coral Reefs of Saurashtra Coast. *Journal of Geology* 83, 1973, fasc. 3, 357–361.

- Haas 1959
F. Haas, Appendix F: Shells Collected by the Expedition. In: H. Field (ed) *An Anthropological Reconnaissance in West Pakistan, 1955. With Appendixes on the Archaeology and Natural History of Baluchistan and Bahawalpur* Papers of the Peabody Museum of Archaeology and Ethnology, Harvard University 52 (Cambridge, Massachusetts 1959) 228.
- Haig 1894
M. R. Haigh, 1894. *The Indus Delta Country, a Memory chiefly on its Ancient Geography and History* (London 1894).
- Hameed-Baloch et al. 2014
A. Hameed-Baloch/M. S. Haneef-Ur-Rehman/ M. A. Kalhoro/M. Aslam Buzdar. *The Curious Case of Mangroves Forest at the Sonmiani bay Area, Lasbela District, Pakistan: A review. Lasbela 2014, University Journal of Science and Technology 3, 2014, 61–74.*
- Hasan 2002
M. U. Hasan, *Baluchistan A Retrospect* (Karachi 2002).
- Hayter 1960
P. J. D. Hayter, *The Ganges and Indus submarine Canyons. Deep Sea Research 6, 1960, 184–186.*
- Hellyer/Aspinal 2006
P. Hellyer/S. Aspinal 2006. *An archaeological and ecological curiosity - Terebralia palustris* (Linnaeus 1767) in the north-east of the Emirate of Abu Dhabi. *Tribulus 16, 2006, 10–13.*
- Hogarth 1999
P. J. Hogarth, *The Biology of Mangroves* (Oxford 1999).
- Holmes 1968
D. A. Holmes, *The Recent History of the Indus. The Geographical Journal 134, 1968, fasc. 3, 367–382.*
- Hughes 1876
A. W. Hughes, *Gazetteer of the Province of Sind, London (2nd edition 1876.)* (reprinted by Indus Publications, Karachi 1996).
- Hughes-Buller 1908
R. Hughes-Buller, *Imperial Gazetteer of India. Provincial Series. Baluchistan* (Calcutta 1908).
- Ibn al-Bitār 1987
Ibn al-Bitār, *Traité des simples. Translated by L. Lucien (1883), Institut du Monde Arabe* (Paris 1987).
- Inam et al. 2007
A. Inam/P. D. Clift/L. Giosan L/A. R. Tabrez/M. Tahir/M. M. Rabbani/M. Danish, *The geographic, geological and oceanographic setting of the Indus River. In: A. Gupta (ed.) Large Rivers: Geomorphology and Management* (London 2007) 333–346.
- Ivory/Lézine 2009
S. J. Ivory/A.-M. Lézine, *Climate and environmental change at the end of the Holocene Humid Period: A pollen record off Pakistan. C. R. Geoscience 341, 2009, 760–769.*
- Jarrige et al. 1995
C. Jarrige/J.-F. Jarrige/R. H. Meadow/G. Quinvron, *Mehrgarh Field Reports 1974-1985 from the Neolithic Times to the Indus Civilization. The Reports of Eleven Seasons of Excavations in Kachi District, Balochistan, by the French Archaeological Mission to Pakistan* (Karachi 1995).
- Jarrige 2004
J.-F. Jarrige, *Le Néolithique des frontières indo-iraniennes: Mehrgarh. In: J. Guilaine (ed.), Aux marges des grands foyers du Néolithique. Périphéries débrutées ou créatrices?* (Paris 2004) 29–60.
- Jarrige/Lechevallier 1980
J.-F. Jarrige/M. Lechevallier, *Les fouilles de Mehrgarh, Pakistan: problèmes chronologique. Paléorient 6, 1980, 253–258.*
- Kathiresan/Rajendran 2005
K. Kathiresan/N. Rajendran, *Mangrove ecosystems of the Indian Ocean Region. Indian Journal of Marine Sciences 34, 2005, fasc. 1, 104–113.*
- Kazi 1999
A. H. Kazi, *The Indus River: Water, Power Resources and Environment. In: A. Meadows/P. Meadows (eds.) The Indus River. Biodiversity, Resources, Humankind* (Karachi 1999) 141–150.
- Kenoyer 2015
J. M. Kenoyer, *The Archaeological Heritage of Pakistan: From the Paleolithic to the Indus Civilization. In: R. D. Long (ed.), A History of Pakistan* (Karachi 2015).
- Kevran 1995
M. Kevran, *Le delta de l'Indus au temps d'Alexandre. Quelques éléments nouveaux pour l'interprétation des sources narratives. Académie des inscriptions et belles-lettres 139:1, 1995, 1, 259–312.*
- Kevran 1999
M. Kevran, *Multiple Ports at the Mouth of the River Indus: Barbarike, Deb, Daybul, Lahori Bandar, Diul Sinde. In: H. P. Ray, (ed.), Archaeology of Seafaring. The Indian Ocean in the Ancient Period* (Delhi 1999) 70–153.
- Khan 1979a
A. R. Khan, *Ancient Settlements in Karachi Region. In: A. Khuhro (ed.), Studies in Geomorphology and Prehistory of Sind, Grassroots 3:2, 1979, 1–24.*
- Khan 1979b
A. R. Khan, *Geomorphology of the Mango Pir Spur. In: A. Khuhro (ed.), Studies in Geomorphology and Prehistory of Sind, Grassroots 3, 1979, fasc. 2 (special issue), 35–46.*
- Khan 1979c
A. R. Khan, *New Archaeological Sites in Las Bela. A Neolithic Settlement Discovered. In A. Khuhro (ed.), Studies in Geomorphology and Prehistory of Sind, Grassroots 3:2, 1979, (special issue) 62–78.*
- Kumaran et al. 2005
K. P. N. Kumaran/K. M. Nair/M. Shindikar/R. B. Limaye/D. Padmalal, *Stratigraphical and palynological appraisal of the Late Quaternary mangrove deposits of the West Coast of India. Quaternary Research 64, 2005, 418–431.*
- Lambrick 1986
H. T. Lambrick, *Sind. A general introduction. History of Sind Series 1* (Hyderabad/Jamshoro 1986) (3rd edition).
- Lechevallier 2003
M. Lechevallier, *L'industrie Lithique de Mehrgarh Fouilles 1974–1985* (Paris 2003).
- Lugo/Snedaker 1974
A. E. Lugo/S. C. Snedaker, *The Ecology of Mangroves. Annual Review of Ecology and Systematics 5, 1974, 39–64.*
- Majumdar 1934
N. C. Majumdar, *Explorations in Sind. Being a report of the exploratory survey carried out during the years*

- 1927-28, 1929-30 and 1930-31. *Memoirs of the Archaeological Survey of India* 48, (Karachi 1934) (1st reprint).
- McCordle 1901
J. W. McCordle, *Ancient India as described in classical literature* (London 1901).
- McCordle 1972
J. W. McCordle, *The Commerce and Navigation of the Erythraean Sea; being a translation of the Periplus Maris Erythraei, by an anonymous writer, and of Arrian's Account of the Voyage of Nearkhos, from the mouth of the Indus to the head of the Persian Gulf followed by Ancient India as described by Ktesias and Knidian being a translation of the abridgement of his "Indika" by Photios and of fragments of that work preserved in other writers* (Amsterdam 1972) (1st reprint).
- McKean 1983
M. B. McKean, *Palynology of Balakot: a Pre-Harappan and Harappan Age site in Las Bela, Pakistan*. Thesis Submitted for PhD Degree (Dallas 1983) (unpublished).
- Meadow 1979
R. H. Meadow, *Prehistoric Subsistence at Balakot: Initial Considerations on the Faunal remains*. In: M. Taddei (ed.), *South Asian Archaeology, Istituto Universitario Orientale. Seminario di Studi Asiatici, Series Minor, VI* (Naples 1977) 275-315.
- Meadows/Meadows 1999
P. S. Meadows/A. Meadows, *The Environmental Impact of the River Indus on the Coastal and Offshore Zones of the Arabian Sea and North-west Indian Ocean*. In: A. Meadows/P. Meadows (eds.) *The Indus River. Biodiversity, Resources, Humankind* (Karachi 1999) 151-171.
- Méry 1996
S. Méry, *Ceramics and Patterns of Exchange across the Arabian Sea and the Persian Gulf in the Early Bronze Age*. In: *Colloquium XXXII. Trade as a Subsistence Strategy. Post Pleistocene Adaptations in Arabia and Early Maritime Trade in the Indian Ocean*. XII International Congress of Prehistoric and Protohistoric Sciences (Forlì 1996) 157-179.
- Méry 2000
S. Méry, *Les céramiques d'Oman et l'Asie moyenne. Une archéologie des échanges à l'Âge du Bronze*. *CRA Monographies* 23 (Paris 2000).
- Minchin 1907
C. F. Minchin, *Las Bela. Text and Appendices* (Karachi 1907).
- Murray 1881
J. A. Murray, *The plants and drugs of Sind, being a systematic account, with descriptions, of the indigenous flora, and notices of the value and uses of their products in commerce, medicine and the arts* (London 1881).
- Naseem et al. 1996
S. Naseem/S. A. Sheikh/M. Qadeeruddin, *Geochemistry and Sedimentology of Jhill Limestone of Gaj Formation, in Cape Monze and Adjoining Area, Karachi*. *Chinese Journal of Geochemistry* 15, 1996, fasc. 3, 213-227.
- Panhwar 1964
M. H. Panhwar, *Ground Water in Hyderabad & Khairpur Divisions* (Hyderabad 1964).
- Panhwar 1983
M. H. Panhwar, *Chronological Dictionary of Sind* (Jamshoro 1983).
- Pathan 1978
M. H. Pathan, *Sind Arab Period. History of Sind Series 3* (Hyderabad 1978) (1st edition).
- Petrie et al. 2010
C. Petrie/F. Khan/R. Knox/K. Thomas/J. Morris, *The investigations of early villages in the hills and on the plains of western South Asia*. In: C. A. Petrie (ed.), *Sheri Khan Tarakai and early village life in the borderlands of north-western Pakistan* (Oxford 2010) 1-28.
- Piggott 1950
S. Piggott, *Prehistoric India to 1000 B.C.* (Harmondsworth 1950).
- Pithawalla 1939
M. B. Pithawalla, *Settlements in the Lower Indus Basin (Sind). Part I. Showing the Influences of Political, Climatic, Geomorphological, Tectonic and Hydrographical Changes in the Region*. *Journal of the Madras Geographical Association* 13:4, 1939, 323-357.
- Pithawalla 1976
M. B. Pithawalla, *A Physical and Economic Geography of Sind (The Lower Indus Basin)* (Hyderabad 1976).
- Plaziat 1995
J.-C. Plaziat, *Modern and fossil mangroves and mangals: their climatic and biogeographic variability*. *Geological Society, London, Special Publications* 83, 1995, 73-96.
- Possehl 2003
G. L. Possehl, *The Indus Civilization. A Contemporary Perspective* (Oxford 2003).
- Qureshi 1990
M. T. Qureshi, *Experimental Plantation for Rehabilitation of Mangrove Forests in Pakistan. First Report. UNDP/UNESCO Regional Project for Research and its Application to the Management of the Mangroves of Asia and the Pacific (RAS/86/120)* (Karachi 1990).
- Rahman 1988
M. Rahman, *Agriculture in Pakistan. Geography of World Agriculture* 13. *Research Institute of Geography, Hungarian Academy of Sciences* (Budapest 1988).
- Rainbird 2007
P. Rainbird, *The Archaeology of Islands* (New York 2010).
- Rasool and Saifullah 2005
F. Rasool/S. M. Saifullah, *A new technique for growing the grey mangrove *Avicennia marina* (Forssk) Vier., in the field*. *Pakistan Journal of Botany* 37:4, 2005, 969-972.
- Ratnagar 2004
S. Ratnagar, *Trading Encounters. From the Euphrates to the Indus in the Bronze Age* (New Delhi 2004).
- Raverty 1895
H. G. Raverty, *The Mihran of Sind and its tributaries: a geographical and historical study* (Calcutta 1895).
- Ray 2003
H. P. Ray, *The Archaeology of Seafaring in Ancient South Asia* (Cambridge 2003).
- Reid et al. 2008
D. G. Reid/P. Dyal/P. Lozouet/M. Glaubrecht/S. T. Williams, *Mudwhelks and mangroves: The evolutionary history of an ecological association (Gastropoda: Potamididae)*. *Molecular Phylogenetics and Evolution* 47, 2008, 680-699.

- Reimer/Reimer 2001
P. J. Reimer/R. W. Reimer, A Marine reservoir correction data base and on-line interface. *Radiocarbon* 43, 2001, fasc. 2A, 461–463.
- Reimer et al. 2013
P. J. Reimer/E. Bard/A. Bayliss/J. W. Beck/P. G. Blackwell/C. Bronk Ramsey/P. M. Grootes/T. P. Guilderson/H. Hafliðason/I. Hajdas/C. Hatté/T. J. Heaton/D. L. Hoffmann/A. G. Hogg/K. F. Hughen/K. F. Kaiser/B. Kromer/S. W. Manning/M. Niu/R. W. Reimer/D. A. Richards/E. M. Scott/J. R. Southon/R. A. Staff/C. S. M. Turney/J. van der Plicht J. 2013. IntCal13 and Marine13 Radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55:4, 2013, 1869–1887.
- Rizwi et al. 1999
S. H. N. Rizwi/M. Nisa/S. M. Haq, Environmental Degradation and Marine Pollution Along the Karachi Coast and Adjacent Creeks in the Indus Delta. In: A. Meadows/P. Meadows (eds.), *The Indus River. Biodiversity, Resources, Humankind* (Karachi 1999) 79–90.
- Rollet 1981
B. Rollet, *Bibliography on mangrove research 1600-1975* (Paris 1981).
- Saifullah et al. 2002
S. M. Saifullah/F. Rasool, 2002. Mangroves of Miani Hor on the North Arabian Sea coast of Pakistan. *Pakistan Journal of Botany* 34, 2002, fasc. 3, 303–310.
- Saliège et al. 2005
J.-F. Saliège/A.-M. Lézine/S. Cleuziou, Estimation de l'effet réservoir ^{14}C marin en mer d'Arabie. *Paléorient* 31:1, 2005, 64–69.
- Sarwar 1992
G. Sarwar, Tectonic setting of the Bela Ophiolites, southern Pakistan. *Tectonophysics* 207, 1992, fasc. 3-4, 359–381.
- Schwadron 2013
M. Schwadron, Prehistoric Shell Landscapes of the Ten Thousand Islands, Florida. In: G. N. Bailey/K. Hardy/A. Camara (eds.), *Shell Energy. Mollusc Shells as Coastal Resources* (Oxford-Oakville 2013) 43–58.
- Shaffer 1986
J. Shaffer, The Archaeology of Baluchistan: A Review. *Newsletter of Baluchistan Studies* 3, 1986, 63–111.
- Shuaib/Tariq Shuaib 1999
S. M. Shuaib/S. M. Tariq Shuaib, Geology and Oil/Gas Presence in the Offshore Indus Basin of Pakistan. In: A. Meadows/P. Meadows (eds.) *The Indus River. Biodiversity, Resources, Humankind* (Karachi 1999) 249–265.
- Siddiqi 1956
M. I. Siddiqi, The Fishermen's Settlements on the Coast of West Pakistan. *Schriften des Geographischen Instituts der Universität Kiel* 16:2, 1956, 1–92.
- Snead 1966
R. E. Snead, *Physical Geography Reconnaissance: Las Bela Coastal Plain, West Pakistan*. Louisiana State University Studies Coastal Studies Series 13 (Baton Rouge 1966) 1–117.
- Snead 1967
R. E. Snead, Recent Morphological Changes along the Coast of West Pakistan. *Annals of the Association of American Geographers* 57:3, 1967, 550–565.
- Snead 1969
R. E. Snead, *Physical Geography Reconnaissance: West Pakistan Coastal Zone*. University of New Mexico Publications in Geography 1. Department of Geography, University of New Mexico (Albuquerque 1969).
- Snead/Frishman 1968
R. E. Snead/S. A. Frishman, Origin of Sands on the East Side of Las Bela Valley, West Pakistan. *Geological Society of America Bulletin* 79, 1968, 1671–1677.
- Snedaker 1984
S. C. Snedaker, Mangroves: A Summary of Knowledge with Emphasis on Pakistan. In: B. U. Haq/J. D. Milliman (eds.) *Marine Geology and Oceanography of the Arabian Sea and Coastal Pakistan* (New York 1984) 254–262.
- Starnini/Biagi 2011
E. Starnini/P. Biagi, The Archaeological Record of the Indus (Harappan) Lithic Production: The Excavation of RH862 Flint Mine and Flint Knapping Workshops on the Rohri Hills (Upper Sindh, Pakistan). *Journal of Asian Civilizations* 34:2, 2011, 1–61.
- Staubwasser et al. 2002
M. Staubwasser/F. Sirocko/P. M. Grootes/H. Herlenkeuser, South Asian monsoon climate change and radiocarbon in the Arabian Sea during early and middle Holocene. *Paleoceanography* 17:4, 2002, 15-1–15-12.
- Staubwasser et al. 2003
M. Staubwasser/F. Sirocko/P. M. Grootes/M. Segl, Climate change at the 4.2 ka BP termination of the Indus valley civilization and Holocene south Asian monsoon variability. *Geophysical Research Letters* 30, 2003, fasc. 8, 7-1–7-4.
- Stein 1943
A. Stein, On Alexander's route into Gedrosia: An Archaeological Tour in Las Bela. *The Geographical Journal* 102:5-6, 1943, 193–226.
- Stewart/Pilkey 1966
R. A. Stewart/O. Pilkey, *Sediments of the Northern Arabian Sea* (Washington D.C 1966).
- Thomas 2015a
K. D. Thomas, Molluscs emergent, Part I: themes and trends in the scientific investigation of mollusc shells as resources for archaeological research. *Journal of Archaeological Sciences* 56, 2015, 133–140. <http://dx.doi.org/10.1016/j.jas.2015.01.024>.
- Thomas 2015b
K. D. Thomas, Molluscs emergent, Part II: themes and trends in the scientific investigation of molluscs and their shells as past human resources. *Journal of Archaeological Sciences* 56, 2015, 159–167. <http://dx.doi.org/10.1016/j.jas.2015.01.015>
- Tomlison 1986
P. B. Tomlison, *The botany of mangroves* (Cambridge 1986).
- Tremenheere 1867
C. W. Tremeneere, On the Lower Portion of the River Indus. *Journal of the Royal Geographical Society of London* 37, 1867, 68–91.
- Uerpmann 1991
H. P. Uerpmann, Radiocarbon dating of shell middens in the Sultanate of Oman. *PACT* 29, 1991, 335–347.
- Uerpmann/Uerpmann 2003
H.-P. Uerpmann/M. Uerpmann, *Stone Age Sites and their Natural Environment. The Capital Area of Northern*

- Oman Part III. Beihefte zum Tübinger Atlas der Vorderen Orients, Reihe A (Naturwissenschaften) Nr. 31/3 (Wiesbaden 2003).
- Vannini et al. 2008
M. Vannini/S. Cannicci/E. Mrabu/R. Rorandelli/F. Frattini, Random walk, zonation and the food searching strategy of *Terebralia palustris* (Mollusca, Potamididae) in Kenya. *Estuarine, Coastal and Shelf Science* 80, 2008, 529–537.
- van Rheede 1678-1703
H. van Rheede, *Hortus Indicus Malabaricus*. Amstelredami 1678-1703, Johannis van Someren et Joannis van Dick (in Latin). <http://dx.doi.org/10.5962/bhl.title.707>.
- von Rad et al. 1999
U. von Rad/M. Schaaf/K. H. Michels/H. Schulz/W. H. Berger/F. Sirocko, A 5000-yr record of climate change in varved sediments from the oxygen minimum zone off Pakistan, northeastern Arabian Sea. *Quaternary Research* 51, 1999, 39–53.
- Vredenburg 1909
E. E. Vredenburg, Report on the Geology of Sarawan, Jhalawan, Mekran and the State of Las Bela, Considered Principally from the Point of View of Economic Development. *Records of the Geological Survey of India* 37, 1909, fasc. 3, 189–215.
- Wells/Coleman 1984
J. T. Wells/J. M. Coleman. 1984. Deltaic Morphology and Sedimentology, with special reference to the Indus River Delta. In: B. U. Haq/J. D. Milliman (eds.) *Marine Geology and Oceanography of the Arabian Sea and Coastal Pakistan* (New York 1984) 85–100.
- Wilhelmy 1968
H. Wilhelmy, Indus delta and Rann of Kutch. *Erdkunde* 23:3, 1968, 177–191.
- Wilhelmy 1986
H. Wilhelmy, The Shifting River. *Sindhological Studies*, Winter 1986, 5–23.
- Wright 2010
R. P. Wright, *The Ancient Indus: Urbanism, Economy and Society* (New York 2010).
- Yoganandan et al. 2013
V. Yoganandan/C. Krishnaiah/K. Selvaraj/G. V. Ravi Prasad/K. Dutta, Monsoonal Fluctuations vs Marine Productivity during Past 10,000 Years - A Study Based on Sediment Core Retrieved from Southeastern Arabian Sea. In: J. Sundaresan/S. Sreekesh/A. L. Ramanathan/L. Sonnenschein/R. Boojh (eds.) *Climate Change and Island and Coastal Vulnerability* (New York 2013) 22–29.
- Zaidi et al. 1999
S. M. S. Zaidi/M. Quadri/G. Hamid/M. Bilal, The Landform Inventory and Genesis in the Mulri Hills Area, Karachi East. *Journal Geographic* 2, 1999, fasc. 1, 39–48.
- Zazzo et al. 2016
A. Zazzo/O. Munoz/E. Badel/I. Béguier/F. Genchi/L. G. Marcucci, A Revised Radiocarbon Chronology of the Aceramic Shell Midden of Ra's Al-Hamra 6 (Muscat, Sultanate of Oman): Implication For Occupational Sequence, Marine Reservoir Age, and Human Mobility, *Radiocarbon*, 2016. Available on CJO 2016 <http://dx.doi.org/10.1017/RDC.2016.3>.
- Zazzo et al. 2012
A. Zazzo/O. Munoz/J.-F. Saliège/C. Moreau, Variability in the marine radiocarbon reservoir effect in Muscat (Sultanate of Oman) during the 4th millennium BC: reflection of taphonomy or environment? *Journal of Archaeological Science* 39, 2012, fasc. 7, 2559–2567.
- Zorzi et al. 2015
C. Zorzi/A. F. Sanchez Goñi/K. Krishnamurthy Anupama/S. Prasad/V. Hanquiez/J. Johnson/L. Giosan, Indian monsoon variations during three contrasting climatic periods: The Holocene, Heinrich Stadial 2 and the last interglacial transition. *Quaternary Science Reviews* 125, 2015, 50–60.
- Амирханов 1997
Х. А. Амирханов, Неолит и Постнеолит Хадрамаута и Махры (Москва 1997).
- Амирханов 2006
Х. А. Амирханов, Каменный век Южной Арабии (Москва 2006).

Paolo Biagi
Department of Asian and North African Studies
Ca' Foscari University
Ca' Cappello, San Polo 2035
I-30125 Venezia, Italia
Email: pavelius@unive.it

Renato Nisbet
Department of Asian and North African Studies
Ca' Foscari University
Ca' Cappello, San Polo 2035
I-30125 Venezia, Italia
Email: renato.nisbet@unive.it

Tiziano Fantuzzi
Department of Humanities
Ca' Foscari University
Palazzo Malcanton-Marcorà
Dorsoduro 3484/D
I-30123 Venezia, Italia
Email: tiziano.fantuzzi@gmail.com

Summary

The research carried out between 2000 and 2014 along the north Arabian Sea coast of Lower Sindh and Las Bela in Balochistan (Pakistan) has shown that the two regions started to be settled, in and around mangroves, during the last three centuries of the 9th millennium BP. The sites discovered during fieldwork are represented by shell middens, shell scatters and fishermen villages, the presence of which was almost unknown until the beginning of the 2000s. Many of the sites were sampled for conventional and AMS radiocarbon dating from mangrove gastropods, and more rarely marine shells. So far 105 sites have been AMS (GrA-) or conventionally (GrN-) radiocarbon-dated, 84 of which from *Terebralia palustris* or *Telescopium*

telescopium adult specimens. This paper describes the results so far obtained from three well-defined macro areas (Lake Siranda, the coastline between Cape Gadani and the Hab River mouth, and the Indus Delta) where this research methodology has been applied. The results contribute to the study of the archaeology of the coastal area of present-day Pakistan, the definition of the sea-shore variations that took place since the middle of the Atlantic period, the disappearance and exploitation of ancient mangroves, and the modes of advance of the Indus Delta up to Historic times. Furthermore it contributes to the study of the early navigation along the northern coast of the Arabian Sea inhabited by different groups of fish-eaters (ichthyophagoi).